

**Military Deployment**  
**Periodic Occupational and Environmental Monitoring Summary (POEMS):**  
**Forward Operating Base (FOB) Ghazni and vicinity, Afghanistan**  
**Calendar Years: (2004 to 2013)**

**AUTHORITY:** This periodic occupational and environmental monitoring summary (POEMS) has been developed in accordance with Department of Defense (DoD) Instructions 6490.03, 6055.05, and JCSM (MCM) 0028-07, See *REFERENCES*.

**PURPOSE:** This POEMS documents the Department of Defense (DoD) assessment of occupational and environmental health (OEH) risk for Forward Operating Base (FOB) Ghazni, its sub-installations, and other base camps in the vicinity that includes Combat Outposts (COP) Band-E Sardeh, Dey Yak, Four Corners, Giro, Kushamond (aka Pacemaker), Qara Bagh (aka Qara Abad), and FOB Vulcan. It presents a qualitative summary of health risks identified at these locations and their potential medical implications. The report is based on information collected from 15 March 2004 through 8 May 2013 to include deployment OEHS sampling and monitoring data (e.g., air, water, and soil), field investigation and health assessment reports, as well as country and area-specific information on endemic diseases.

This assessment assumes that environmental sampling at FOB Ghazni and vicinity during this period was performed at representative exposure points selected to characterize health risks at the *population-level*. Due to the nature of environmental sampling, the data upon which this report is based may not be fully representative of all the fluctuations in environmental quality or capture unique occurrences. While one might expect health risks pertaining to historic or future conditions at this site to be similar to those described in this report, the health risk assessment is limited to 15 March 2004 through 8 May 2013.

The POEMS can be useful to inform healthcare providers and others of environmental conditions experienced by individuals deployed to FOB Ghazni and vicinity during the period of this assessment. However, it does not represent an individual exposure profile. Individual exposures depend on many variables such as; how long, how often, where and what someone is doing while working and/or spending time outside. Individual outdoor activities and associated routes of exposure are extremely variable and cannot be identified from or during environmental sampling. Individuals who sought medical treatment related to OEH exposures while deployed should have exposure/treatment noted in their medical record on a Standard Form (SF) 600 (Chronological Record of Medical Care).

**SITE DESCRIPTION:**

The base camps are located in the Ghazni Province and neighboring Paktika Province for COP Kushamond. The city of Ghazni serves as the provincial capital of Ghazni province. It lies on the Kabul-Kandahar Highway, and has historically functioned as an important trade center. A large water reservoir, created by the Band E Sardeh Dam, is located near COP Band-E Sardeh and the border of Paktika province. Ghazni and Paktika provinces are primarily rural agricultural societies. The region is mainly hilly and interspersed with seasonal river valleys. The terrain gains elevation and becomes more rugged to the north.

**SUMMARY:** Conditions that may pose a Moderate or greater health risk are summarized in Table 1. Table 2 provides population based risk estimates for identified OEH conditions at FOB Ghazni and vicinity. As indicated in the detailed sections that follow Table 2, controls established to reduce health risk were factored into this assessment. In some cases, e.g. ambient air, specific controls are noted, but not routinely available/feasible.

**Table 1: Summary of Occupational and Environmental Conditions with MODERATE or Greater Health Risk**

**Short-term health risks & medical implications:**

The following hazards may be associated with potential acute health effects in some personnel during deployment at Forward Operating Base (FOB) Ghazni and vicinity that includes Combat Outpost (COP) Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni, COP Giro, COP Kushamond, COP Qarabagh, and FOB Vulcan:

Food/waterborne diseases (e.g., bacterial diarrhea, hepatitis A, typhoid/paratyphoid fever, diarrhea-cholera, diarrhea- protozoal, brucellosis, hepatitis E); other endemic diseases (malaria, cutaneous leishmaniasis (acute), Crimean-Congo hemorrhagic fever, sandfly fever, typhus-miteborne, leptospirosis, Tuberculosis (TB), rabies, anthrax, Q fever); heat stress; and cold stress. For food/waterborne diseases (e.g., bacterial diarrhea, hepatitis A, typhoid/paratyphoid fever, diarrhea- cholera, diarrhea- protozoal, brucellosis, hepatitis E), if ingesting local food and water, the health effects can temporarily incapacitate personnel (diarrhea) or result in prolonged illness (hepatitis A, typhoid/paratyphoid fever, brucellosis, hepatitis E). Risks from food/waterborne diseases may have been reduced with preventive medicine controls and mitigation, which includes hepatitis A and typhoid fever vaccinations and only drinking from approved water sources in accordance with standing CENTCOM policy. For other vector-borne endemic diseases (malaria, cutaneous leishmaniasis (acute), Crimean- Congo hemorrhagic fever, sandfly fever, typhus-miteborne), these diseases may constitute a significant risk due to exposure to biting vectors; risk reduced to 'Low' by proper wear of the treated uniform, application of repellent to exposed skin, bed net use, and appropriate chemoprophylaxis, as well as minimizing areas of standing water and other vector- breeding areas. For water contact diseases (leptospirosis) activities involving extensive contact with surface water increase risk. For respiratory diseases (TB), personnel in close-quarter conditions could have been at risk for person-to-person spread. Animal contact diseases (rabies, anthrax, Q fever), pose year-round risk. For heat stress, risk can be greater during months of June through September, and greater for susceptible persons including those older than 45, of low fitness level, unacclimatized, or with underlying medical conditions, and those under operational constraints (equipment, PPE, vehicles). Risks from heat stress may have been reduced with preventive medicine controls, work-rest cycles, proper hydration and nutrition, and mitigation. For cold stress, the risk of Non-Freezing Cold Injuries (NFCI), such as chilblain, trench foot, and hypothermia, is greater during the months of November through March. Risks from cold stress may have been reduced with protective measures such as use of the buddy system, limiting exposure during cold weather, proper hydration and nutrition, and proper wear of issued protective clothing.

Air quality: For inhalable coarse particulate matter less than 10 micrometers in diameter (PM<sub>10</sub>), the PM<sub>10</sub> overall short-term risk was not evaluated due to insufficient data. For inhalable fine particulate matter less than 2.5 micrometers in diameter (PM<sub>2.5</sub>), the PM<sub>2.5</sub> overall short-term risk was not evaluated due to no available data. However, exposures to PM<sub>10</sub> and PM<sub>2.5</sub> may vary, as conditions may vary, and may result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel while at this site, particularly exposures to high levels of dust such as during high winds or dust storms. For PM<sub>10</sub> and PM<sub>2.5</sub>, certain subgroups of the deployed forces (e.g., those with pre-existing asthma/cardio-pulmonary conditions) are at greatest risk of developing notable health effects. For burn pits, although the short-term risk for PM<sub>10</sub> and for PM<sub>2.5</sub> was not evaluated due to no data available for PM<sub>10</sub> and insufficient data available for PM<sub>2.5</sub>, there were operating burn pits utilized at or around COP Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni, COP Giro, COP Kushamond, and COP Qarabagh – see Sections 10.2 and 10.7. For burn pits, exposures may vary, and exposure to high levels of PM<sub>10</sub> and to PM<sub>2.5</sub> in the smoke may also result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel and certain subgroups while at this site. Although most short-term health effects from exposure to particulate matter and burn pit smoke should have resolved post- deployment, providers should be prepared to consider the relationship between deployment exposures and current complaints. Some individuals may have sought treatment for acute respiratory irritation during their time at FOB Bostick and COP Monti and vicinity. Personnel who reported with symptoms or required treatment while at this site should have exposure and treatment noted in medical record (e.g., electronic medical record and/or on a Standard Form (SF) 600 (*Chronological Record of Medical Care*)).

**Long-term health risks & medical implications:**

The following hazards may be associated with potential chronic health effects in some personnel during deployment at FOB Ghazni and vicinity that includes (COP) Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni, COP Giro, COP Kushamond, COP Qarabagh, and FOB Vulcan:

Air quality: For inhalable fine particulate matter less than 2.5 micrometers in diameter (PM<sub>2.5</sub>), the overall long-term risk was not evaluated due to no available data. Inhalable coarse particulate matter less than 10 micrometers in diameter (PM<sub>10</sub>) was not evaluated for long-term risk due to no available health guidelines. However, the area was a dusty desert environment, and conditions may have varied. In addition, for burn pits, although the long-term risk for PM<sub>10</sub> and for PM<sub>2.5</sub> was not evaluated due to no available data for PM<sub>10</sub> and insufficient data available for PM<sub>2.5</sub>, there were documented operating burn pits utilized at or around at COP Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni, COP Giro, COP Kushamond, and COP Qarabagh, and conditions may have varied – see Sections 10.2 and 10.7. For inhalational exposure to high levels of dust, PM<sub>10</sub> and PM<sub>2.5</sub>, such as during high winds or dust storms, and for exposure to burn pit smoke, it is considered possible that some otherwise healthy personnel, who were exposed for a long-term period to dust and

particulate matter, could develop certain health conditions (e.g., reduced lung function, cardiopulmonary disease). Personnel with a history of asthma or cardiopulmonary disease could potentially be more likely to develop such chronic health conditions. While the dust and particulate matter exposures and exposures to burn pits are acknowledged, at this time there were no specific recommended, post-deployment medical surveillance evaluations or treatments. Providers should still consider overall individual health status (e.g., any underlying conditions/susceptibilities) and any potential unique individual exposures (such as burn pits/barrels, incinerators, occupational or specific personal dosimeter data) when assessing individual concerns. Certain individuals may need to be followed/evaluated for specific occupational exposures/injuries (e.g., annual audiograms as part of the medical surveillance for those enrolled in the Hearing Conservation Program; and personnel covered by Respiratory Protection Program and/or Hazardous Waste/Emergency Responders Medical Surveillance).

**Table 2. Population-Based Health Risk Estimates - FOB Ghazni and vicinity that includes COP Band-E Sardeh, COP Dey Yak, COP Four Corners, COP Giro, COP Kushamond, COP Qarabagh, and FOB Vulcan<sup>1, 2</sup>**

| Source of Identified Health Risk <sup>3</sup>                                 | Unmitigated Health Risk Estimate <sup>4</sup>  | Control Measures Implemented  | Residual Health Risk Estimate <sup>4</sup>   |
|---|--|---|--|
| <b>AIR</b>  |  |   |  |
| Particulate matter less than 10 micrometers in diameter (PM <sub>10</sub> )   | Short-term: Insufficient data were available for risk analysis. Daily levels vary, acute health effects (e.g., upper respiratory tract irritation) more pronounced during peak days. More serious effects are possible in susceptible persons (e.g., those with asthma/existing respiratory diseases).   | Limiting strenuous physical activities when air quality is especially poor; and actions such as closing tent flaps, windows, and doors.   | Short-term: Insufficient data were available for risk analysis. Daily levels vary, acute health effects (e.g., upper respiratory tract irritation) more pronounced during peak days. More serious effects are possible in susceptible persons (e.g., those with asthma/existing respiratory diseases).                         |
|   | Long-term: No health guidelines  |   | Long-term: No health guidelines  |
| Particulate matter less than 2.5 micrometers in diameter (PM <sub>2.5</sub> ) | Short-term: No data were available for risk analysis. A majority of the time mild acute (short term) health effects are anticipated; certain peak levels may produce mild eye, nose, or throat irritation in some personnel and pre-existing health conditions (e.g., asthma, or cardiopulmonary diseases) may be exacerbated.                       | Limiting strenuous physical activities when air quality is especially poor; and actions such as closing tent flaps, windows, and doors.   | Short-term: No data were available for risk analysis. A majority of the time mild acute (short term) health effects are anticipated; certain peak levels may produce mild eye, nose, or throat irritation in some personnel and pre-existing health conditions (e.g., asthma, or cardiopulmonary diseases) may be exacerbated. |
|   | Long-term: No data were available for risk analysis. A small percentage of personnel may be at increased risk for developing chronic conditions. Particularly those more susceptible to acute effects (e.g., those with asthma/existing respiratory diseases).   |   | Long-term: No data were available for risk analysis. A small percentage of personnel may be at increased risk for developing chronic conditions. Particularly those more susceptible to acute effects (e.g., those with asthma/existing respiratory diseases).   |
| <b>ENDEMIC DISEASE</b>  |  |   |  |
| Food borne/Waterborne (e.g., diarrhea-bacteriological)                        | Short-term: Variable; High (bacterial diarrhea, hepatitis A, typhoid fever) to Moderate (diarrhea-cholera, diarrhea- protozoal, brucellosis, hepatitis E) if ingesting local food/water, the health effects can temporarily incapacitate personnel (diarrhea) or result in prolonged illness (hepatitis A, typhoid fever, hepatitis E, brucellosis). | Preventive measures include hepatitis A and typhoid fever vaccination and consumption of food and water only from approved sources.   | Short-term: Low to none  |
|   | Long-term: none identified   |   | Long-term: No data available   |
| Arthropod Vector Borne  | Short-term: Variable; High for malaria, Moderate for leishmaniasis - cutaneous (acute), Crimean-Congo hemorrhagic fever, sandfly fever, typhus-miteborne; and Low for, the plague and West Nile fever.   | Preventive measures include proper wear of treated uniform, application of repellent to exposed skin, bed net use, minimizing areas of standing water and appropriate chemoprophylaxis. | Short-term: Low  |
|   | Long-term: Low for Leishmaniasis-visceral infection.   |   | Long-term: No data available   |
| Water-Contact (e.g. wading, swimming)   | Short-term: Moderate for leptospirosis   | Recreational swimming in surface waters not likely in this area of Afghanistan during this time period.   | Short-term: Low for leptospirosis.   |
|   | Long-term: No data available   |   | Long-term: No data available   |
| Respiratory   | Short-term: Variable; Moderate for tuberculosis (TB) to Low for meningococcal meningitis.  | Providing adequate living and work space; medical screening; vaccination.   | Short-term: Low  |

|                                   |  |   |  |
|-----------------------------------|--|---|--|
|                                   | Long-term: No data available   |   | Long-term: No data available   |
| Animal Contact                    | Short-term: Variable; Moderate for rabies, anthrax, Q-fever to Low for H5N1 avian influenza.   | Prohibiting contact with, adoption, or feeding of feral animals IAW U.S. Central Command (CENTCOM) General Order (GO) 1B. Risks are further reduced in the event of assessed contact by prompt post-exposure rabies prophylaxis IAW The Center for Disease Control's (CDC) Advisory Committee on Immunization Practices guidance. | Short-term: No data available  |
|                                   | Long-term: Low (Rabies)  |   | Long-term: No data available   |
| <b>VENOMOUS ANIMAL/ INSECTS</b>   |  |   |  |
| Snakes, scorpions, and spiders    | Short-term: Low; If encountered, effects of venom vary with species from mild localized swelling (e.g. <i>Scorpiops lindbergi</i> ) to potentially lethal effects (e.g. <i>Gloydus halys</i> ).  | Risk reduced by avoiding contact, proper wear of uniform (especially footwear), and proper and timely treatment.  | Short-term: Low; If encountered, effects of venom vary with species from mild localized swelling (e.g. <i>Scorpiops lindbergi</i> ) to potentially lethal effects (e.g. <i>Gloydus halys</i> ).  |
|                                   | Long-term: No data available   |   | Long-term: No data available   |
| <b>HEAT/COLD STRESS</b>           |  |   |  |
| Heat                              | Short-term: Low to Moderate; The unmitigated risk of heat injury in unacclimatized or susceptible personnel, and those under operational constraints is Low from October – May, Moderate for September, and high from June to August based on historical temperature data. | Work-rest cycles, proper hydration and nutrition, and Wet Bulb Globe Temperature (WBGT) monitoring.   | Short-term: The risk of heat injury was reduced to low. However, the risk may be greater to unacclimatized or susceptible populations (older, previous history of heat injury, poor physical condition, underlying medical/health conditions), and those under operational constraints (equipment, PPE, vehicles). |
|                                   | Long-term: Low, The long-term risk was Low. However, the risk may be greater to certain susceptible persons—those older (i.e., greater than 45 years), in lesser physical shape, or with underlying medical/health conditions.   |   | Long-term: Low, The long-term risk is Low. However, the risk may be greater to certain susceptible persons—those older (i.e., greater than 45 years), in lesser physical shape, or with underlying medical/health conditions.  |
| Cold                              | Short-term: The risk of Non-Freezing Cold Injuries (NFCI), such as chilblain, trench foot, and hypothermia, is Moderate from November – March based on historical temperature and precipitation data.  | Risks from cold stress reduced with protective measures such as use of the buddy system, limiting exposure during cold weather, proper hydration and nutrition, and proper wear of issued protective clothing.  | Short-term: Low risk of cold stress/injury.  |
|                                   | Long-term: Low; Long-term health implications from cold injuries are rare but can occur, especially from more serious injuries such as frost bite.   |   | Long-term: Low; Long-term health implications from cold injuries are rare but can occur, especially from more serious injuries such as frost bite.   |
| <b>Unique Incidents/ Concerns</b> |  |   |  |
| Burn Pits                         | Short-term: No data were available for PM <sub>10</sub> and insufficient data were available for PM <sub>2.5</sub> at burn pits located at COP Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni,   | Control measures may have included locating burn pits downwind of prevailing winds, increased distance from   | Short-term: No data were available for PM <sub>10</sub> and insufficient data were available for PM <sub>2.5</sub> at burn pits located at COP Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni,   |

|  |  |   |  |
|--|--|---|--|
|  | <p>COP Giro, COP Kushamond, and COP Qarabagh. Burn pit exposures may vary, and exposure to high levels of PM<sub>10</sub> and PM<sub>2.5</sub> in the smoke may also result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel and certain subgroups while at this site.</p>  | <p>living and working areas when possible, and improved waste segregation and management techniques</p> | <p>COP Giro, COP Kushamond, and COP Qarabagh. Burn pit exposures may vary, and exposure to high levels of PM<sub>10</sub> and PM<sub>2.5</sub> in the smoke may also result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel and certain subgroups while at this site.</p>  |
|  | <p>Long-term: No data were available for PM<sub>10</sub> and insufficient data were available for PM<sub>2.5</sub> at COP Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni, COP Giro, COP Kushamond, and COP Qarabagh. Burn pit exposures may vary, and exposure to high levels of PM<sub>10</sub> and PM<sub>2.5</sub> in the smoke may be associated with some otherwise healthy personnel, who were exposed for a long-term period, could develop certain health conditions (e.g., reduced lung function, cardiopulmonary disease). Personnel with a history of asthma or cardiopulmonary disease could potentially be more likely to develop such chronic health conditions.</p> |   | <p>Long-term: No data were available for PM<sub>10</sub> and insufficient data were available for PM<sub>2.5</sub> at COP Band-E Sardeh, COP Dey Yak, COP Four Corners, FOB Ghazni, COP Giro, COP Kushamond, and COP Qarabagh. Burn pit exposures may vary, and exposure to high levels of PM<sub>10</sub> and PM<sub>2.5</sub> in the smoke may be associated with some otherwise healthy personnel, who were exposed for a long-term period, could develop certain health conditions (e.g., reduced lung function, cardiopulmonary disease). Personnel with a history of asthma or cardiopulmonary disease could potentially be more likely to develop such chronic health conditions.</p> |

<sup>1</sup>This Summary Table provides a qualitative estimate of population-based short- and long-term health risks associated with the occupational environment conditions at FOB Ghazni and vicinity. It does not represent an individual exposure profile. Actual individual exposures and health effects depend on many variables. For example, while a chemical may have been present in the environment, if a person did not inhale, ingest, or contact a specific dose of the chemical for adequate duration and frequency, then there may have been no health risk. Alternatively, a person at a specific location may have experienced a unique exposure which could result in a significant individual exposure. Any such person seeking medical care should have their specific exposure documented in an SF600.

<sup>2</sup> This assessment is based on specific environmental sampling data and reports obtained from 15 March 2004 through 8 May 2013. Sampling locations are assumed to be representative of exposure points for the camp population but may not reflect all the fluctuations in environmental quality or capture unique exposure incidents.

<sup>3</sup>This Summary Table is organized by major categories of identified sources of health risk. It only lists those sub-categories specifically identified and addressed at FOB Ghazni and vicinity. The health risks are presented as Low, Moderate, High or Extremely High for both acute and chronic health effects. The health risk level is based on an assessment of both the potential severity of the health effects that could be caused and probability of the exposure that would produce such health effects. Details can be obtained from the USAPHC/Army Institute of Public Health (AIPH). Where applicable, "None Identified" is used when though a potential exposure is identified, no health risks of either a specific acute or chronic health effects are determined. More detailed descriptions of OEH exposures that are evaluated but determined to pose no health risk are discussed in the following sections of this report.

<sup>4</sup>Health risks in this Summary Table are based on quantitative surveillance thresholds (e.g. endemic disease rates; host/vector/pathogen surveillance) or screening levels, e.g. Military Exposure Guidelines (MEGs) for chemicals. Some previous assessment reports may provide slightly inconsistent health risk estimates because quantitative criteria such as MEGs may have changed since the samples were originally evaluated and/or because this assessment makes use of all historic site data while previous reports may have only been based on a select few samples.

## 1 Discussion of Health Risks at FOB Ghazni and vicinity, Afghanistan by Source

The following sections provide additional information about the OEH conditions summarized above. All risk assessments were performed using the methodology described in the U.S. Army Public Health Command Technical Guide 230, *Environmental Health Risk Assessment and Chemical Exposure Guidelines for Deployed Military Personnel* (USAPHC TG230, reference 9). All OEH risk estimates represent residual risk after accounting for preventive controls in place. Occupational exposures and exposures to endemic diseases are greatly reduced by preventive measures. For environmental exposures related to airborne dust, there are limited preventive measures available, and available measures have little efficacy in reducing exposure to ambient conditions.

## 2 Air

### 2.1 Site-Specific Sources Identified

FOB Ghazni and vicinity are situated in a dusty semi-arid desert environment. Inhalational exposure to high levels of dust and particulate matter, such as during high winds or dust storms, may result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel. Additionally, certain subgroups of the deployed forces (e.g., those with pre-existing asthma/cardio pulmonary conditions) are at greatest risk of developing notable health effects.

### 2.2 Particulate matter

Particulate matter (PM) is a complex mixture of extremely small particles suspended in the air. The PM includes solid particles and liquid droplets emitted directly into the air by sources such as: power plants, motor vehicles, aircraft, generators, construction activities, fires, and natural windblown dust. The PM can include sand, soil, metals, volatile organic compounds (VOC), allergens, and other compounds such as nitrates or sulfates that are formed by condensation or transformation of combustion exhaust. The PM composition and particle size vary considerably depending on the source. Generally, PM of health concern is divided into two fractions: PM<sub>10</sub>, which includes coarse particles with a diameter of 10 micrometers or less, and fine particles less than 2.5 micrometers (PM<sub>2.5</sub>), which can reach the deepest regions of the lungs when inhaled. Exposure to excessive PM is linked to a variety of potential health effects.

### 2.3 Particulate matter, less than 10 micrometers (PM<sub>10</sub>)

#### 2.3.1 Exposure Guidelines:

|  |  |
|--|--|
| Short Term (24-hour) PM <sub>10</sub> (micrograms per cubic meter, $\mu\text{g}/\text{m}^3$ ):                                 | Long-term PM <sub>10</sub> MEG ( $\mu\text{g}/\text{m}^3$ ):                     |
| <ul style="list-style-type: none"> <li>Negligible MEG = 250</li> <li>Marginal MEG = 420</li> <li>Critical MEG = 600</li> </ul> | <ul style="list-style-type: none"> <li>Not defined and not available.</li> </ul> |

#### 2.3.2 Sample data/Notes:

FOB Ghazni: A total of 4 valid PM<sub>10</sub> air samples were collected from 22 June 2006 to 7 May 2013. The range of 24-hour PM<sub>10</sub> concentrations was 31  $\mu\text{g}/\text{m}^3$  – 314  $\mu\text{g}/\text{m}^3$ , with an average concentration of 199  $\mu\text{g}/\text{m}^3$ . COP Kushamond (aka Pacemaker): A total of 3 valid PM<sub>10</sub> air samples were collected from 31 March 2011 to 6 May 2011. The range of 24-hour PM<sub>10</sub> concentrations was 67  $\mu\text{g}/\text{m}^3$  – 447  $\mu\text{g}/\text{m}^3$  with an average concentration of 207  $\mu\text{g}/\text{m}^3$ .

### 2.3.3 Short-term health risks:

**Not Evaluated.** There were insufficient data to characterize the short-term PM<sub>10</sub> health risk for FOB Ghazni and COP Kushamond.

### 2.3.4 Long-term health risk:

**Not Evaluated-no available health guidelines.** The U.S. Environmental Protection Agency (EPA) has retracted its long-term standard (national ambient air quality standards, NAAQS) for PM<sub>10</sub> due to an inability to clearly link chronic health effects with chronic PM<sub>10</sub> exposure levels.

## 2.4 Particulate Matter, less than 2.5 micrometers (PM<sub>2.5</sub>)

### 2.4.1 Exposure Guidelines:

Short Term (24-hour) PM<sub>2.5</sub> (µg/m<sup>3</sup>):

- Negligible MEG = 65
- Marginal MEG = 250
- Critical MEG = 500

Long-term (1year) PM<sub>2.5</sub> MEGs (µg/m<sup>3</sup>):

- Negligible MEG = 15
- Marginal MEG = 65.

### 2.4.2 Sample data/Notes:

FOB Ghazni: A single valid PM<sub>2.5</sub> air sample was collected on 8 May 2013. The 24-hour PM<sub>2.5</sub> concentration was 29 µg/m<sup>3</sup>.

COP Kushamond (aka Pacemaker): A single valid PM<sub>2.5</sub> air sample was collected on 9 July 2012. The 24-hour PM<sub>2.5</sub> concentration was 41 µg/m<sup>3</sup>.

### 2.4.3 Short-term and Long-term health risks:

**Not evaluated.** There were insufficient data to characterize the short-term PM<sub>2.5</sub> health risk for FOB Ghazni and COP Kushamond.

## 2.5 Airborne Metals

### 2.5.1 Exposure Guidelines:

No airborne metals were detected above short- or long-term MEGs.

### 2.5.2 Sample data/Notes:

FOB Ghazni: A total of 4 valid PM<sub>10</sub> airborne metal samples were collected from 31 March 2011 to 6 May 2011. None of the detected metal concentrations exceeded the short- or long-term MEGs.

COP Kushamond: A total of 3 valid PM<sub>10</sub> airborne metal samples were collected from 31 March 2011 to 6 May 2011. None of the detected metal concentrations exceeded the short- or long-term MEGs.

### 2.5.3 Short- and long-term health risks:

None identified based on the available sampling data.

## 2.6 Volatile Organic Compounds (VOC)

### 2.6.1 Exposure Guidelines:

No airborne metals were detected above short- or long-term MEGs.

### 2.6.2 Sample data/Notes:

No valid volatile organic chemical (VOC) air samples were available for analysis.

### 2.6.3 Short and long-term health risks:

**Not Evaluated.**

## 3 Soil

### 3.1 Site-Specific Sources Identified

#### 3.2 Sample data/Notes:

The primary soil contamination exposure pathways are dermal contact and dust inhalation. Typical parameters analyzed for included semi volatile organic compounds (SVOCs), heavy metals, polychlorinated biphenyls (PCBs), pesticides, herbicides. If the contaminant was known or suspected, other parameters may have been analyzed for (i.e., total petroleum hydrocarbons (TPH) and polycyclic aromatic hydrocarbons (PAH) near fuel spills). For the risk assessment, personnel are assumed to remain at this location for 6 months to 1 year.

COP Band-E Sardeh: A total of 3 valid surface soil samples were collected on 12 July 2012 to assess OEH health risk to deployed personnel. The percent of the population exposed to soil and associated dust in the sampled areas was from 50% - 75%. No chemicals were detected above the 1 year negligible MEG.

COP Dey Yak: A total of 6 valid surface soil samples were collected on 11 April 2011 to assess OEH health risk to deployed personnel. The percent of the population exposed to soil and associated dust in the sampled areas was > 75% for 3 samples, from 50% - 75% for 2 samples, and from 10% - 25% for 1 sample. No chemicals were detected above the 1 year negligible MEG in any samples.

COP Four Corners: A total of 3 valid surface soil samples were collected on 16 July 2012 to assess OEH health risk to deployed personnel. The percent of the population exposed to soil and associated dust in the sampled areas was 50 < 75%. No chemicals were detected above the 1 year negligible MEG in any samples.

FOB Ghazni: A total of 11 valid surface soil samples were collected from 25 September 2006 to 7 May 2013 to assess OEH health risk to deployed personnel. The percent of the population exposed to soil and associated dust in the sampled areas was > 75% for 3 samples, between 50 < 75% for 3 samples, between 25 < 50% for 3 samples and < 10% for 2 samples. No chemicals were detected above the 1 year negligible MEG.

COP Giro: A total of 3 valid surface soil samples were collected on 13 June 2008 to assess OEH health risk to deployed personnel. The percent of the population exposed to soil and associated dust in the sampled areas was > 75%. No chemicals were detected above the 1 year negligible MEG.

COP Kushamond (aka Pacemaker): A total of 16 valid surface soil samples were collected from 27 October 2007 to 10 July 2012 to assess OEH health risk to deployed personnel. The percent of the population exposed to soil and associated dust in the sampled areas was > 75% for 7 samples, between 50 < 75% for 3 samples, between 25 < 50% for 5 samples and between 10 < 25% for 1 sample. No chemicals were detected above the 1 year negligible MEG.

COP Qarabagh: A single valid surface soil sample was collected on 14 June 2012 to assess OEH health risk to deployed personnel. The percent of the population exposed to soil and associated dust in the sampled areas was > 75%. No chemicals were detected above the 1 year negligible MEG.

### 3.3 Short-term health risk:

**Not an identified source of health risk.** Currently, sampling data for soil are not evaluated for short term (acute) health risks.

### 3.4 Long-term health risk:

**None identified based on available sample data.** No parameters exceeded 1-year Negligible MEGs.

## 4 Water

In order to assess the health risk to U.S. personnel from exposure to water in theater, the USAPHC identified the most probable exposure pathways. These are based on the administrative information provided on the field data sheets submitted with the samples taken over the time period being evaluated. It is assumed that 100% of all U.S. personnel at FOB Ghazni and vicinity will be directly exposed to reverse osmosis water purification unit (ROWPU) treated, disinfected fresh bulk water, and untreated water since this classification of water is primarily used for personal hygiene, showering, cooking, and for use at vehicle wash racks. Field data sheets indicate that bottled water is the only approved source of drinking water.

### 4.1 Drinking Water: Bottled or Packaged Water

#### 4.1.1 Site-Specific Sources Identified

#### 4.1.2 Sample data/Notes:

No valid bottled water samples were available for analysis.

#### 4.1.3 Short-term and long-term health risk:

**Not Evaluated.**

### 4.2 Non-Drinking Water:

#### 4.2.1 Site-Specific Sources Identified

Although the primary route of exposure for most microorganisms is ingestion of contaminated water, dermal exposure to some microorganisms, chemicals, and biologicals may also cause adverse health effects. Complete exposure pathways would include drinking, brushing teeth, personal hygiene, cooking, providing medical and dental care using a contaminated water supply or during dermal contact at vehicle or aircraft wash racks.

#### 4.2.2 Sample data/Notes:

To assess the potential for adverse health effects to troops the following assumptions were made about dose and duration: All U.S. personnel at this location were expected to remain at this site for approximately 1 year. A conservative (protective) assumption is that personnel routinely consumed less than 5L/day of non-drinking water for up to 365 days (1-year). It is further assumed that control measures and/or personal protective equipment were not used.

COP Band-E Sardeh: A single disinfected bulk water (Non-Drinking) sample and 2 untreated bulk water (Non-Drinking) samples collected from 24 September 2010 to 12 July 2012 were evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

COP Dey Yak: A single disinfected bulk water (Non-Drinking) sample and 2 untreated bulk water (Non-Drinking) samples collected from 11 April 2011 to 27 June 2012 were evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

COP Four Corners: A single disinfected bulk water (Non-Drinking) sample and 2 untreated bulk water (Non-Drinking) samples collected from 23 September 2010 to 16 July 2012 were evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

FOB Ghazni: A total of 7 disinfected bulk water (Non-Drinking) samples and 19 untreated bulk water (Non-Drinking) samples collected from 26 June 2006 to 7 May 2013 were evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

COP Giro: A total of 2 untreated bulk water (Non-Drinking) samples collected on 13 June 2006 and 16 June 2012 were evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

COP Kushamond: A total of 3 untreated bulk water (Non-Drinking) samples and 3 untreated bulk water (Non-Drinking) samples collected from 16 August 2009 to 29 January 2013 were evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

COP Four Corners: A single untreated bulk water (Non-Drinking) sample collected on 14 June 2012 was evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

FOB Vulcan: A single untreated bulk water (Non-Drinking) sample collected on 7 November 2007 was evaluated for this health risk assessment. No chemicals were detected at levels above the short or long-term MEGs.

#### 4.2.3 Short and long-term health risks:

**None identified based on available sample data.** All collected samples were below the short and long-term Negligible MEGs.

## 5 Military Unique

### 5.1 Chemical Biological, Radiological Nuclear (CBRN) Weapons

No specific hazard sources were documented in the Defense Occupational and Environmental Health Readiness System (DOEHRS), or the Military Exposure Surveillance Library (MESL) from 15 March 2004 to 8 February 2013 timeframe.

## 5.2 Depleted Uranium (DU)

No specific hazard sources were documented in the DOEHRS, or MESL from 15 March 2004 to 8 February 2013 timeframe.

## 5.3 Ionizing Radiation

No specific hazard sources were documented in the DOEHRS, or MESL from 15 March 2004 to 8 February 2013 timeframe.

## 5.4 Non-Ionizing Radiation

No specific hazard sources were documented in the DOEHRS, or MESL from 15 March 2004 to 8 February 2013 timeframe.

# 6 Endemic Diseases

This document lists the endemic diseases reported in the region, its specific health risks and severity and general health information about the diseases. CENTCOM Modification (MOD) 11 (Reference 11) lists deployment requirements, to include immunizations and chemoprophylaxis, in effect during the timeframe of this POEMS.

## 6.1 Foodborne and Waterborne Diseases

Food borne and waterborne diseases in the area are transmitted through the consumption of local food and water. Local unapproved food and water sources (including ice) are heavily contaminated with pathogenic bacteria, parasites, and viruses to which most U.S. Service Members have little or no natural immunity. Effective host nation disease surveillance does not exist within the country. Only a small fraction of diseases are identified or reported in host nation personnel. Diarrheal diseases are expected to temporarily incapacitate a very high percentage of U.S. personnel within days if local food, water, or ice is consumed. Hepatitis A and typhoid fever infections typically cause prolonged illness in a smaller percentage of unvaccinated personnel. Vaccinations are required for DoD personnel and contractors. In addition, although not specifically assessed in this document, significant outbreaks of viral gastroenteritis (e.g., norovirus) and food poisoning (e.g., *Bacillus cereus*, *Clostridium perfringens*, *Staphylococcus*) may occur. Key disease risks are summarized below:

Mitigation strategies were in place and included consuming food and water from approved sources, vaccinations (when available), frequent hand washing and general sanitation practices.

FOB Ghazni: Total coliform was repeatedly found in water samples according to Base Camp Assessments (BCA) from 3 October 2004 to 30 June 2006.

### 6.1.1 Diarrheal diseases (bacteriological)

**High, mitigated to Low:** Diarrheal diseases are expected to temporarily incapacitate a very high percentage of personnel (potentially over 50% per month) within days if local food, water, or ice is consumed. Field conditions (including lack of hand washing and primitive sanitation) may facilitate person-to-person spread and epidemics. Typically mild disease treated in outpatient setting; recovery and return to duty in less than 72 hours with appropriate therapy. A small proportion of infections may require greater than 72 hours limited duty, or hospitalization.

### 6.1.2 Hepatitis A, typhoid/paratyphoid fever, and diarrhea-protozoal

**High, mitigated to Low:** Unmitigated health risk to U.S. personnel is high year round for hepatitis A and typhoid/paratyphoid fever, and Moderate for diarrhea-protozoal. Mitigation was in place to reduce the risks to low. Hepatitis A, typhoid/paratyphoid fever, and diarrhea-protozoal disease may cause prolonged illness in a small percentage of personnel (less than 1% per month). Although much rarer, other potential diseases in this area that are also considered a Moderate risk include: hepatitis E, diarrhea-cholera, and brucellosis.

#### 6.1.3 Short-term Health Risks:

**Low:** The overall unmitigated short-term risk associated with food borne and waterborne diseases are considered High (bacterial diarrhea, hepatitis A, typhoid/paratyphoid fever) to Moderate (diarrhea-cholera, diarrhea-protozoal, brucellosis) to Low (hepatitis E) if local food or water is consumed. Preventive Medicine measures reduced the risk to Low. Confidence in the health risk estimate is high.

#### 6.1.4 Long-term Health Risks:

**None identified based on available data.**

## 6.2 Arthropod Vector-Borne Diseases

During the warmer months, the climate and ecological habitat support populations of arthropod vectors, including mosquitoes, ticks, mites, and sandflies. Significant disease transmission is sustained countrywide, including urban areas. Malaria, the major vector-borne health risk in Afghanistan, is capable of debilitating a high percentage of personnel for up to a week or more. Mitigation strategies were in place and included proper wear of treated uniforms, application of repellent to exposed skin, and use of bed nets and chemoprophylaxis (when applicable). Additional methods included the use of pesticides, reduction of pest/breeding habitats, and engineering controls.

### 6.2.1 Malaria

**High, mitigated to Low** Potential unmitigated risk to U.S. personnel is High during warmer months (typically April through November) but reduced to low with mitigation measures. Malaria incidents are often associated with the presence of agriculture activity, including irrigation systems and standing water, which provide breeding habitats for vectors. A small number of cases may occur among personnel exposed to mosquito (*Anopheles* spp.) bites. Malaria incidents may cause debilitating febrile illness typically requiring 1 to 7 days of inpatient care, followed by return to duty. Severe cases may require intensive care or prolonged convalescence.

### 6.2.2 Leishmaniasis

**Moderate, mitigated to Low:** The disease risk is Moderate during the warmer months when sandflies are most prevalent, but reduced to low with mitigation measures. Leishmaniasis is transmitted by sand flies. There are two forms of the disease; cutaneous (acute form) and visceral (a more latent form of the disease). The leishmaniasis parasites may survive for years in infected individuals and this infection may go unrecognized by physicians in the U.S. when infections become symptomatic years later. Cutaneous infection is unlikely to be debilitating, though lesions may be disfiguring. Visceral leishmaniasis disease can cause severe febrile illness which typically requires hospitalization with convalescence over 7 days.

### 6.2.3 Crimean-Congo hemorrhagic fever

**Moderate, mitigated to Low:** Unmitigated risk is moderate, but reduced to low with mitigation measures. Crimean-Congo hemorrhagic fever occurs in rare cases (less than 0.1% per month attack rate in indigenous personnel) and is transmitted by tick bites or occupational contact with blood or secretions from infected animals. The disease typically requires intensive care with fatality rates from 5% to 50%.

### 6.2.4 Sandfly fever

**Moderate, mitigated to Low:** Sandfly fever has a Moderate risk with potential disease rates from 1% to 10% per month under worst case conditions. Mitigation measures reduced the risk to low. The disease is transmitted by sandflies and occurs more commonly in children though adults are still at risk. Sandfly fever disease typically resulted in debilitating febrile illness requiring 1 to 7 days of supportive care followed by return to duty.

### 6.2.5 Plague

**Low:** Potential health risk to U.S. personnel is Low year round. Bubonic plague typically occurred as sporadic cases among people who come in contact with wild rodents and their fleas during work, hunting, or camping activities. Outbreaks of human plague are rare and typically occur in crowded urban settings associated with large increases in infected commensal rats (*Rattus rattus*) and their flea populations. Some untreated cases of bubonic plague may develop into secondary pneumonic plague. Respiratory transmission of pneumonic plague is rare but has the potential to cause significant outbreaks. Close contact is usually required for transmission. In situations where respiratory transmission of plague is suspected, weaponized agent must be considered. Extremely rare cases (less than 0.01% per month attack rate) could occur. Incidence could result in potentially severe illness which may require more than 7 days of hospitalization and convalescence.

### 6.2.6 Typhus-miteborne (scrub typhus)

**Moderate, mitigated to Low:** Potential health risk to U.S. personnel is Moderate during warmer months (typically March through November) when vector activity is highest. Mitigation measures reduced the risk to low. Mite-borne typhus is a significant cause of febrile illness in local populations with rural exposures in areas where the disease is endemic. Large outbreaks have occurred when non-indigenous personnel such as military forces enter areas with established local transmission. The disease is transmitted by the larval stage of trombiculid mites (chiggers), which are typically found in areas of grassy or scrubby vegetation, often in areas which have undergone clearing and regrowth. Habitats may include sandy beaches, mountain deserts, cultivated rice fields, and rain forests. Although data are insufficient to assess potential disease rates, attack rates can be very high (over 50%) in groups of personnel exposed to heavily infected "mite islands" in focal areas. The disease can cause debilitating febrile illness typically requiring 1 to 7 days of inpatient care, followed by return to duty.

### 6.2.7 West Nile fever

**Low:** West Nile fever is present. The disease is maintained by the bird population and transmitted to humans via mosquito vector. Typically, infections in young, healthy adults were asymptomatic although fever, headache, tiredness, body aches (occasionally with a skin rash on trunk of body), and swollen lymph glands can occur. This disease is associated with a low risk estimate.

### 6.2.9 Short-term health risks:

**Low:** The unmitigated health risk estimate is High for malaria (infection rate of less than 1% per month), Moderate for leishmaniasis-cutaneous (acute), Crimean-Congo hemorrhagic fever, sandfly fever, typhus-miteborne; and Low for, the plague and West Nile fever. Health risk is reduced to low by proper wear of the uniform, application of repellent to exposed skin, and appropriate chemoprophylaxis. Confidence in health risk estimate was high.

### 6.2.10 Long-term health risks:

**Low:** The unmitigated risk is moderate for leishmaniasis-visceral (chronic). Risk is reduced to Low by proper wear of the uniform and application of repellent to exposed skin. Confidence in the risk estimate is high.

## 6.3 Water Contact Diseases

Operations or activities that involve extensive water contact may result in personnel being temporarily debilitated with leptospirosis in some locations. Leptospirosis health risk typically increases during flooding. In addition, although not specifically assessed in this document, bodies of surface water are likely to be contaminated with human and animal waste. Activities such as wading or swimming may result in exposures to enteric diseases such as diarrhea and hepatitis via incidental ingestion of water. Prolonged water contact also may lead to the development of a variety of potentially debilitating skin conditions such as bacterial or fungal dermatitis. Mitigation strategies were in place and included avoiding water contact and recreational water activities, proper wear of uniform (especially footwear), and protective coverings for cuts/abraded skin.

### 6.3.1 Leptospirosis

**Moderate, mitigated to Low:** Human infections occur seasonally (typically April through November) through exposure to water or soil contaminated by infected animals and is associated with wading, and swimming in contaminated, untreated open water. The occurrence of flooding after heavy rainfall facilitates the spread of the organism because as water saturates the environment leptospirosis present in the soil passes directly into surface waters. Leptospirosis can enter the body through cut or abraded skin, mucous membranes, and conjunctivae. Infection may also occur from ingestion of contaminated water. The acute, generalized illness associated with infection may mimic other tropical diseases (for example, dengue fever, malaria, and typhus), and common symptoms include fever, chills, myalgia, nausea, diarrhea, cough, and conjunctival suffusion. Manifestations of severe disease can include jaundice, renal failure, hemorrhage, pneumonitis, and hemodynamic collapse. Recreational activities involving extensive water contact may result in personnel being temporarily debilitated with leptospirosis. Incidence could result in debilitating febrile illness typically requiring 1 to 7 days of inpatient care, followed by return to duty; some cases may require prolonged convalescence. This disease is associated with a Moderate health risk estimate.

### 6.3.2 Short-term health risks:

**Low:** Unmitigated Health risk of leptospirosis is Moderate during warmer months. Mitigation measures reduce the risk to Low. Confidence in the health risk estimate is high.

### 6.3.3 Long-term health risks:

**None identified based on available data.**

## 6.4 Respiratory Diseases

Although not specifically assessed in this document, deployed U.S. forces may be exposed to a wide variety of common respiratory infections in the local population. These include influenza, pertussis, viral upper respiratory infections, viral and bacterial pneumonia, and others. The U.S. military populations living in close-quarter conditions are at risk for substantial person-to-person spread of respiratory pathogens. Influenza is of particular concern because of its ability to debilitate large numbers of unvaccinated personnel for several days. Mitigation strategies were in place and included routine medical screenings, vaccination, enforcing minimum space allocation in housing units, implementing head-to-toe sleeping in crowded housing units, implementation of proper personal protective equipment (PPE) when necessary for healthcare providers and detention facility personnel.

### 6.4.1 Tuberculosis (TB)

**Moderate, mitigated to Low:** Potential health risk to U.S. personnel is Moderate, mitigated to Low, year round. Transmission typically requires close and prolonged contact with an active case of pulmonary or laryngeal TB, although it also can occur with more incidental contact. The Army Surgeon General has defined increased risk in deployed Soldiers as indoor exposure to locals or third country nationals of greater than one hour per week in a highly endemic active TB region. Additional mitigation included active case isolation in negative pressure rooms, where available.

### 6.4.2 Meningococcal meningitis

**Low:** Meningococcal meningitis poses a Low risk and is transmitted from person to person through droplets of respiratory or throat secretions. Close and prolonged contact facilitates the spread of this disease. Meningococcal meningitis is potentially a very severe disease typically requiring intensive care; fatalities may occur in 5-15% of cases.

### 6.4.3 Short-term health risks:

**Low:** Moderate (TB) to Low (for meningococcal meningitis). Overall risk was reduced to Low with mitigation measures. Confidence in the health risk estimate is high.

### 6.4.4 Long-term health risks:

**None identified based on available data.** Tuberculosis is evaluated as part of the post deployment health assessment (PDHA). A TB skin test is required post-deployment if potentially exposed and is based upon individual service policies.

## 6.5 Animal-Contact Diseases

### 6.5.1 Rabies

**Moderate, mitigated to Low:** Rabies posed a year-round moderate risk. Occurrence in local animals was well above U.S. levels due to the lack of organized control programs. Dogs are the primary reservoir of rabies in Afghanistan, and a frequent source of human exposure. Rabies is transmitted by exposure to the virus-laden saliva of an infected animal, typically through bites, but could occur from scratches contaminated with the saliva. A U.S. Army Soldier stationed in Afghanistan died of rabies on 31 August 2011 (Reference 12). Laboratory results indicated the Soldier was infected from contact with a dog while deployed. Although the vast majority (>99%) of persons who develop rabies disease will do so within a year after an exposure, there have been rare reports of individuals presenting with rabies disease up to six years or more after their last known risk exposure. Mitigation strategies included

command emphasis of CENTCOM GO 1B, reduction of animal habitats, active pest management programs, and timely treatment of feral animal scratches/bites.

#### 6.5.2 Anthrax

**Low:** Anthrax cases are rare in indigenous personnel, and pose a Low risk to U.S. personnel. Anthrax is a naturally occurring infection; cutaneous anthrax is transmitted by direct contact with infected animals or carcasses, including hides. Eating undercooked infected meat may result in contracting gastrointestinal anthrax. Pulmonary anthrax is contracted through inhalation of spores and is extremely rare. Mitigation measures included consuming approved food sources, proper food preparation and cooking temperatures, avoidance of animals and farms, dust abatement when working in these areas, vaccinations, and proper PPE for personnel working with animals.

#### 6.5.3 Q-Fever

**Moderate, mitigated to Low:** Potential health risk to U.S. personnel is Moderate, but mitigated to Low, year round. Rare cases are possible among personnel exposed to aerosols from infected animals, with clusters of cases possible in some situations. Significant outbreaks (affecting 1-50%) can occur in personnel with heavy exposure to barnyards or other areas where animals are kept. Unpasteurized milk may also transmit infection. The primary route of exposure is respiratory, with an infectious dose as low as a single organism. Incidence could result in debilitating febrile illness, sometimes presenting as pneumonia, typically requiring 1 to 7 days of inpatient care followed by return to duty. Mitigation strategies in place as listed in paragraph 6.5.2 except for vaccinations.

#### 6.5.4 H5N1 avian influenza

**Low:** Potential health risk to U.S. personnel is Low. Although H5N1 avian influenza (AI) is easily transmitted among birds, bird-to-human transmission is extremely inefficient. Human-to-human transmission appears to be exceedingly rare, even with relatively close contact. Extremely rare cases (less than 0.01% per month attack rate) could occur. Incidence could result in very severe illness with fatality rate higher than 50 percent in symptomatic cases. Mitigation strategies included avoidance of birds/poultry and proper cooking temperatures for poultry products.

#### 6.5.5 Short-term health risks:

**Low:** The short-term unmitigated risk is Moderate for rabies, and Q-fever, to Low for anthrax, and H5N1 avian influenza. Mitigation measures reduced the overall risk to Low. Confidence in risk estimate is high.

#### 6.5.6 Long-term health risks:

**Low:** A Low long term risk exists for rabies because, in rare cases, the incubation period for rabies can be several years.

## 7 Venomous Animals

All information was taken directly from the Clinical Toxinology Resources web site from the University of Adelaide, Australia (Reference 2). The species listed below have home ranges that overlap the location of FOB Ghazni and vicinity, and may present a health risk if they are encountered by personnel. See Section 9 for more information about pesticides and pest control measures.

## 7.1 Spiders

- *Latrodectus dahlia* (widow spider): Severe envenoming possible, potentially lethal. However, venom effects are mostly minor and even significant envenoming is unlikely to be lethal.

## 7.2 Scorpions

- *Androctonus amoreuxi*, and *Androctonus baluchicus*: Severe envenoming possible, potentially lethal. Severe envenoming may produce direct or indirect cardio toxicity, with cardiac arrhythmias, cardiac failure. Hypovolaemic hypotension possible in severe cases due to fluid loss through vomiting and sweating.
- *Buthacus striffleri*, *Compsobuthus afghanus*, *Compsobuthus rugosulus*, *Mesobuthus caucasicus*, *Mesobuthus eupeus*, *Mesobuthus macmahoni*, *Orthochirus afghanus*, *Orthochirus bicolor*, *Orthochirus. Jalalabadensis*, *Orthochirus pallidus*, *Orthochirus samrchelsis*, and *Orthochirus scrobiculosus*: There are a number of dangerous Buthid scorpions, but there are also some known to cause minimal effects only. Without clinical data it is unclear where these species fit within that spectrum.
- *Hottentotta alticola*, and *Hottentotta saulcyi*: Moderate envenoming possible but unlikely to prove lethal. Stings by these scorpions are likely to cause only short lived local effects, such as pain, without systemic effects.
- *Scorpiops afghanus*: Mild envenoming only, not likely to prove lethal. Stings by these scorpions are likely to cause only short lived local effects, such as pain, without systemic effects.

## 7.3 Snakes

- *Boiga trigonata* (Common Cat Snake), and *Telescopus rhinopoma* (leopard viper): Unlikely to cause significant envenoming; Bites by these rear fanged Colubrid snakes are rarely reported. They are likely to cause minimal to moderate local effects and no systemic effects.
- *Echis multisquamatus* (central Asian saw-scaled viper), *Echis sochureki* (Sochurek's saw-scaled viper), *Gloydius halys* (Haly's Pit Viper): Severe envenoming possible, potentially lethal. Bites may cause moderate to severe coagulopathy and haemorrhagins causing extensive bleeding.
- *Macrovipera lebetina obtuse* (Levantine Viper): Severe envenoming possible, potentially lethal. Bites may cause mild to severe local effects, shock & coagulopathy.
- *Naja oxiana* (Oxus cobra): Severe envenoming possible, potentially lethal. Bites can cause systemic effects, principally flaccid paralysis.
- *Platyceps rhodorachis* (Jan's desert racer): Mild envenoming only, not likely to prove lethal. Requires symptomatic treatment only.

## 7.4 Short-term health risk:

**Low:** If encountered, effects of venom vary with species from mild localized swelling (e.g. widow spider) to potentially lethal effects (e.g., Haly's Pit Viper). See effects of venom above. Mitigation strategies included avoiding contact, proper wear of uniform (especially footwear), and timely medical treatment. Confidence in the health risk estimate is low (Reference 9, Table 3-6).

## 7.5 Long-term health risk:

**None identified.**

## 8 Heat/Cold Stress

### 8.1 Heat

Summer (June - September) monthly mean daily maximum temperatures range from 79.7 degrees Fahrenheit (°F) to 87.4 °F with an average temperature of 69.7 °F based on historical climatological data from the U.S. Air Force Combat Climatology Center, 14<sup>th</sup> Weather Squadron. The health risk of heat stress/injury based on temperatures alone is Low (< 78 °F) from October – May, Moderate (78-81.9°F) for September, and high (82-87.9°F) from June to August. However, work intensity and clothing/equipment worn pose greater health risk of heat stress/injury than environmental factors alone (Reference 6). Managing risk of hot weather operations included monitoring work/rest periods, proper hydration, and taking individual risk factors (e.g. acclimation, weight, and physical conditioning) into consideration. Risk of heat stress/injury was reduced with preventive measures

## 8.1.1 Short-term health risk:

**Low to Moderate, mitigated to Low:** The risk of heat injury was reduced to low through preventive measures such as work/rest cycles, proper hydration and nutrition, and monitoring wet bulb globe temperature (WBGT). Risk of heat injury in unacclimatized or susceptible populations (older, previous history of heat injury, poor physical condition, underlying medical/health conditions), and those under operational constraints (equipment, PPE, vehicles) is Low. Confidence in the health risk estimate is low (Reference 9, Table 3-6).

## 8.1.2 Long-term health risk:

**Low:** The long-term risk is Low. However, the risk may be greater for certain susceptible persons—those older (i.e., greater than 45 years), in lesser physical shape, or with underlying medical/health conditions. Long-term health implications from heat injuries are rare but may occur, especially from more serious injuries such as heat stroke. It is possible that high heat in conjunction with various chemical exposures may increase long-term health risks, though specific scientific evidence is not conclusive. Confidence in these risk estimates is medium (Reference 9, Table 3-6).

### 8.2 Cold

## 8.2.1 Short-term health risks:

Winter (December - March) mean daily minimum temperatures range from 12.9 °F to 29.3 °F with an average temperature of 28.3 °F based on historical climatological data from the U.S. Air Force Combat Climatology Center, 14<sup>th</sup> Weather Squadron. Because even on warm days a significant drop in temperature after sunset by as much as 40 °F can occur, there is a risk of cold stress/injury from November – March.

The risk assessment for Non-Freezing Cold Injuries (NFCI), such as chilblain, trench foot, and hypothermia, is Moderate based on historical temperature and precipitation data. Frostbite is possible when temperatures drop below freezing. Personnel may encounter significantly lower temperatures during field operations at higher altitudes. As with heat stress/injuries, cold stress/injuries are largely dependent on operational and individual factors instead of environmental factors alone.

**Low to Moderate, mitigated to Low:** The risk of cold injury was reduced to low through preventive measures such as work/rest cycles, proper hydration and nutrition, and monitoring Wet Bulb Globe Temperature (WBGT). Confidence in the health risk estimate is low.

#### 8.1.2 Long-term health risk:

**Low:** The health risk of cold injury is Low. Confidence in the health risk estimate is high.

## 9 Noise

### 9.1 Continuous

No specific hazard sources were documented in the DOEHRS or MESL from 15 March 2004 to 8 February 2013 timeframe.

#### 9.1.1 Short and long-term health risks:

**Not evaluated**

### 9.2 Impulse

No specific hazard sources were documented in the DOEHRS or MESL from 15 March 2004 to 8 February 2013 timeframe.

#### 9.2.1 Short-term and Long-term health risks:

**Not evaluated.**

## 10 Unique Incidents/Concerns

### 10.1 Potential environmental contamination sources

DoD personnel are exposed to various chemical, physical, ergonomic, and biological hazards in the course of performing their mission. These types of hazards depend on the mission of the unit and the operations and tasks which the personnel are required to perform to complete their mission. The health risk associated with these hazards depends on a number of elements including what materials are used, how long the exposure last, what is done to the material, the environment where the task or operation is performed, and what controls are used. The hazards can include exposures to heavy metal particulates (e.g. lead, cadmium, manganese, chromium, and iron oxide), solvents, fuels, oils, and gases (e.g. carbon monoxide, carbon dioxide, oxides of nitrogen, and oxides of sulfur). Most of these exposures occur when performing maintenance task such as painting, grinding, welding, engine repair, or movement through contaminated areas. Exposures to these occupational hazards can occur through inhalation (air), skin contact, or ingestion; however exposures through air are generally associated with the highest health risk.

### 10.2 Waste Sites/Waste Disposal

FOB Ghazni: According to the USFOR-A Environmental Database, open burning of solid waste was performed on FOB Ghazni. The OEHSA dated 5/7/13 indicates that there were two solid waste incinerators and one medical waste incinerator on the base. The solid waste incinerators were not operating during the date of the OEHSA.

COP Band-E Sardeh: A burn pit was used for disposal of solid waste. Waste water was hauled off site (BCA Band-E Sardeh 2012).

COP Dey Yak: An offsite burn pit was used for disposal of solid waste. Waste water was hauled off site (BCA Dey Yak 2012).

COP Four Corners: A burn pit was used for disposal of solid waste.

COP Giro: A burn pit and incinerator were used for disposal of solid waste (BCA Giro 2011). Waste water was pumped off site.

COP Kushamond: An offsite burn pit was used for disposal of solid waste. However, the distance from the burn pit to COP Kushamond is not known.

COP Qarabagh: An offsite burn pit was used for disposal of solid waste. Waste water was hauled off site (BCA 2012).

10.2.1 Short and long-term health risks:

**Undetermined**

### 10.3 Fuel/petroleum products/industrial chemical spills

COP Qarabagh: Signs of fuel spillage around distribution system and within/outside of secondary containment (BCA 2012).

10.3.1 Short and long-term health risks:

**Undetermined**

### 10.4 Pesticides/Pest Control:

The health risk of exposure to pesticide residues is considered within the framework of typical residential exposure scenarios, based on the types of equipment, techniques, and pesticide products that have been employed, such as enclosed bait stations for rodenticides, various handheld equipment for spot treatments of insecticides and herbicides, and a number of ready-to-use (RTU) methods such as aerosol cans and baits. The control of rodents required the majority of pest management inputs, with the acutely toxic rodenticides staged as solid formulation lethal baits placed in tamper-resistant bait stations indoors and outdoors throughout cantonment areas. Nuisance insects, including biting and stinging insects such as bees, wasps, and ants, also required significant pest management inputs. Use of pesticides targeting against these pests generally involved selection of compounds with low mammalian toxicity and short-term residual using pinpoint rather than broadcast application techniques. A total of 144 monthly pesticide application reports in the MESL data portal for COP Band-E Sardeh, COP Four Corners, FOB Ghazni, COP Giro, and COP Kushamond (13 June 2007 to 4 January 2013) list the usage of pesticides on the site.

For each pesticide product applied during this period, the EPA approved label has been archived, providing a framework how each pesticide handled and applied (see below).

#### 10.4.1 Rodenticides

Polybutylenes, polyisobutylenes, pyrethrins, diphacinone, resmethrin, and brodifacoum were used to control rodents.

#### 10.4.2 Insecticides

Insecticides used to control ants, cockroaches, flies, mosquitoes, and wasps include: *hydramethylnon*, *β-Cyfluthrin*, *Imidacloprid*, Lambda-cyhalothrin, Pyrethrins, Piperonyl Butoxide, Orthoboric Acid, Fipronil, Z- 9- Tricosene, DEET, Glyphosate, Methomyl, *Bacillus thuringiensis* subspecies *israelensis*, (S)-Methoprene, and n-Octyl bicycloheptene dicarboximide.

#### 10.4.3 Short-term and Long-term health risks

**Not evaluated.**

### 10.5 Asbestos

No specific hazard sources were documented in the DOEHS or MESL from 15 March 2004 to 8 February 2013 timeframe for all the bases except COP Four Corners. Possible asbestos containing material was observed within the roofing material on the MWR Building and in the cover of a former ammo bunker in two OEHSAs performed in 2012. No samples were collected to verify if the material did indeed contain asbestos.

#### 10.5.1 Short-term and Long-term health risks:

**Not evaluated.**

### 10.6 Lead Based Paint

No specific hazard sources were documented in the DOEHS or MESL from 15 March 2004 to 8 February 2013 timeframe.

#### 10.6.1 Short-term and Long-term health risks:

**Not evaluated.**

### 10.7 Burn Pit

No information on the presence/absence of burn pit(s) or incinerator(s) were found for FOB Vulcan.

COP Band-E Sardeh: A burn pit for disposal of solid waste was located on the opposite side of the Afghanistan National Army compound. There were no incinerators on site (BCA Band-E Sardeh 2012).

COP Dey Yak: Local national used two-burn pits offsite with a third at some point not visible from perimeter wall. Both observed pits were 10 m x 15 m with depth of 1.2 m. There were no incinerators on site (BCA Dey Yak 2012).

COP Four Corners: A burn pit for disposal of solid waste was located in the corner of COP Four Corners adjacent to HESCO barrier. Solders were exposed to smoke from burning operations when the wind blows smoke into the COP.

FOB Ghazni: According to the USFOR-A Environmental Database, open burning of solid waste was performed on FOB Ghazni. OEHSA dated 5/7/13 indicates that there were two solid waste incinerators and one medical waste incinerator on the base. The solid waste incinerators were not operating during the date of the OEHSA.

COP Giro: A burn pit adjacent to COP was used for disposal of solid waste. An incinerator was used for burning plastics and metals (BCA Giro 2011). Note, the BCA conducted on 15-16 June 2012 indicates that no incinerators were in use; however burn out latrines were used.

COP Kushamond: A burn pit located outside the perimeter was used for disposal of solid waste.

COP Qarabagh: A burn pit located outside the perimeter was used for disposal of solid waste. The burn Pit dimensions 6m x 4.5 m x 2m operated by local national. Local national was observed placing trash on edge of pit rather than inside it.

While not specific to FOB Ghazni and vicinity, the consolidated epidemiological and environmental sampling and studies on burn pits that have been conducted as of the date of this publication have been unable to determine whether an association does or does not exist between exposures to emissions from the burn pits and long-term health effects (Reference 7). The Institute of Medicine committee's review of the literature and the data suggests that service in Iraq or Afghanistan (i.e., a broader consideration of air pollution than exposure only to burn pit emissions) may be associated with long-term health effects, particularly in susceptible (e.g., those who have asthma) or highly exposed subpopulations, such as those who worked at the burn pit. Such health effects would be due mainly to high ambient concentrations of PM from both natural and anthropogenic sources, including military sources. If that broader exposure to air pollution turns out to be relevant, potentially related health effects of concern are respiratory and cardiovascular effects and cancer. Susceptibility to the PM health effects could be exacerbated by other exposures, such as stress, smoking, local climatic conditions, and co-exposures to other chemicals that affect the same biologic or chemical processes. Individually, the chemicals measured at burn pit sites in the study were generally below concentrations of health concern for general populations in the United States. However, the possibility of exposure to mixtures of the chemicals raises the potential for health outcomes associated with cumulative exposure to combinations of the constituents of burn pit emissions and emissions from other sources.

#### 10.7.1 Particulate Matter, less than 2.5 micrometers (PM<sub>2.5</sub>)

##### 10.7.1.2 Exposure Guidelines:

###### Short Term (24-hour) PM<sub>2.5</sub> (µg/m<sup>3</sup>):

- Negligible MEG = 65
- Marginal MEG = 250
- Critical MEG = 500

###### Long-term (1year) PM<sub>2.5</sub> MEGs (µg/m<sup>3</sup>):

- Negligible MEG = 15
- Marginal MEG = 65

##### 10.7.1.3 Sample data/Notes:

COP Giro: A single valid PM<sub>2.5</sub> air sample was collected on 15 June 2012. The 24-hour PM<sub>2.5</sub> concentration was 27 µg/m<sup>3</sup>.

COP Qarabagh: A single valid PM<sub>2.5</sub> air sample was collected on 13 June 2012. The 24-hour PM<sub>2.5</sub> concentration was 177 µg/m<sup>3</sup>.

10.7.1.4 Short- and long-term health risks:

**Not Evaluated.** There were insufficient data to characterize the short- and long-term PM<sub>2.5</sub> health risk for COP Giro, and COP Qarabagh.

## 11 References<sup>1</sup>

1. Casarett and Doull's Toxicology: the Basic Science of Exposures, Chapter 2- Principles of Toxicology; Fifth Edition, McGraw Hill, New York.
2. Clinical Toxinology Resources: <http://www.toxinology.com/>. University of Adelaide, Australia.
3. Defense Occupational and Environmental Health Readiness System (referred to as the DOEHRSEH database) at <https://doehrs-ih.csd.disa.mil/Doehrs/>. Department of Defense (DoD) Instruction 6490.03, *Deployment Health*, 2006.
4. DoDI 6055.05, Occupational and Environmental Health, 2008.
5. DoD MESL Data Portal: <https://mesl.apgea.army.mil/mesl/>. Some of the data and reports used may be classified or otherwise have some restricted distribution.
6. Goldman RF. 2001. Introduction to heat-related problems in military operations. *In*: Textbook of military medicine: medical aspects of harsh environments Vol. 1, Pandolf KB, and Burr RE (Eds.), Office of the Surgeon General, Department of the Army, Washington DC.
7. IOM (Institute of Medicine). 2011. Long-term health consequences of exposure to burn pits in Iraq and Afghanistan. Washington, DC: The National Academies Press.
8. Joint Staff Memorandum (MCM) 0028-07, Procedures for Deployment Health Surveillance, 2007.
9. USA PHC TG230, June 2013 Revision.
10. USACHPPM. 2008. Particulate Matter Factsheet; 64-009-0708, 2008.
11. Modification 11 to United States Central Command Individual Protection and Individual Unit Deployment Policy, 2 December 2011.
12. CDC. 2012. Morbidity and Mortality Weekly Report. Imported Human Rabies in a U.S. Army Soldier. May 4, 2012. 61(17); 302-305.

---

<sup>1</sup> NOTE. The data are currently assessed using the 2013 TG230. The general method involves an initial review of the data which eliminates all chemical substances not detected above 1-yr negligible MEGs. Those substances screened out are not considered acute or chronic health hazards so are not assessed further. For remaining substances, acute and chronic health effects are evaluated separately for air water (soil is only evaluated for long term risk). This is performed by deriving separate short-term and long term population exposure level and estimates (referred to as population exposure point concentrations (PEPC)) that are compared to MEGs derived for similar exposure durations. If less than or equal to negligible MEG the risk is Low. If levels are higher than negligible then there is a chemical-specific toxicity and exposure evaluation by appropriate SMEs, which includes comparison to any available marginal, critical or catastrophic MEGs. For drinking water 15 L/day MEGs are used for the screening while site specific 5-15 L/day are used for more detailed assessment. For nondrinking water (such as that used for personal hygiene or cooking) the 'consumption rate' is limited to 2 L/day (similar to the EPA) which is derived by multiplying the 5 L/day MEG by a factor of 2.5. This value is used to conservatively assess non drinking uses of water.

## 12 Where Do I Get More Information?

If a provider feels that the Service member's or Veteran's current medical condition may be attributed to specific OEH exposures at this deployment location, he/she can contact the Service-specific organization below. Organizations external to DoD should contact DoD Force Health Protection and Readiness (FHP & R).

**Army Institute of Public Health** Phone: (800) 222-9698. <http://phc.amedd.army.mil/>

**Navy and Marine Corps Public Health Center (NMCPHC)** (formerly NEHC) Phone: (757) 953-0700. [www.nmcpbc.med.navy.mil](http://www.nmcpbc.med.navy.mil)

**U.S. Air Force School of Aerospace Medicine (USAFSAM)** (formerly AFIOH) Phone: (888) 232-3764. <http://www.wpafb.af.mil/afri/711hpw/usafsam.asp>

**DoD Force Health Protection and Readiness (FHP & R)** Phone: (800) 497-6261. <http://fhp.osd.mil>