

Military Deployment
Periodic Occupational and Environmental Monitoring Summary (POEMS):
Camp Panther, Afghanistan
Calendar Year: 2008

AUTHORITY: This periodic occupational and environmental monitoring summary (POEMS) has been developed in accordance with Department of Defense (DoD) Instructions 6490.03, 6055.05, and JCSM (MCM) 0017-12 (References 2-4).

PURPOSE: This POEMS documents the Department of Defense (DoD) assessment of occupational and environmental health (OEH) risk for Camp Panther, Afghanistan. It presents a qualitative summary of OEH risks identified at this location and their potential medical implications. The report is based on information collected in 2008 to include deployment OEH surveillance sampling and monitoring data (e.g., air, water, and soil), field investigation and health assessment reports, as well as country and area-specific information on endemic diseases.

This assessment assumes that environmental sampling at Camp Panther during this period was performed at representative exposure points selected to characterize health risks at the *population-level*. Due to the nature of environmental sampling, the data upon which this report is based may not be fully representative of all the fluctuations in environmental quality or capture unique occurrences. While one might expect health risks pertaining to historic or future conditions at this site to be similar to those described in this report, the health risk assessment is limited to 2008.

The POEMS can be useful to inform healthcare providers and others of environmental conditions experienced by individuals deployed to Camp Panther during the period of this assessment. However, it does not represent an individual exposure profile. Individual exposures depend on many variables such as; how long, how often, where and what someone is doing while working and/or spending time outside. Individual outdoor activities and associated routes of exposure are extremely variable and cannot be identified from or during environmental sampling. Individuals who sought medical treatment related to OEH exposures while deployed should have exposure/treatment noted in their medical record on a Standard Form (SF) 600 (Chronological Record of Medical Care).

SITE DESCRIPTION: Due to the fact that Camp Panther was a new camp being built in Afghanistan in 2008 and troops were not likely stationed at the Camp for long, very little information exists on the location of the Camp within Afghanistan. Further inquiry with the COCOM did not yield any additional information on the location. Likewise, information in the DOEHRS does not identify a specific location. Therefore, all risk assessment is general for Afghanistan (with the exception of specific samples taken) and are not specific for Camp Panther.

SUMMARY: Conditions that may pose a Moderate or greater health risk are summarized in Table 1. Table 2 provides population based risk estimates for identified OEH conditions at Camp Panther. As indicated in the detailed sections that follow Table 2, controls established to reduce health risk were factored into this assessment. In some cases, e.g., ambient air, specific controls are noted, but not routinely available/feasible.

The appearance of external hyperlinks does not constitute endorsement by the United States Department of Defense (DoD) of the linked websites, or the information, products or services contained therein. The DoD does not exercise any editorial, security, or other control over the information you may find at these locations.

CLEARED
For Open Publication

10
Mar 22, 2021

Department of Defense
OFFICE OF PREPUBLICATION AND SECURITY REVIEW

Table 1: Summary of Occupational and Environmental Conditions with MODERATE or Greater Health Risk

Short-term health risks & medical implications:

The following hazards may be associated with potential acute health effects in some personnel during deployment at Camp Panther:

Food/waterborne diseases (e.g., bacterial diarrhea, hepatitis A, typhoid/paratyphoid fever, diarrhea-cholera, diarrhea-protozoal, brucellosis, hepatitis E); other endemic diseases (malaria, Crimean-Congo hemorrhagic fever, leishmaniasis-cutaneous (acute), sandfly fever, typhus-miteborne (scrub typhus), leptospirosis, Tuberculosis (TB), rabies, Q fever, soil transmitted helminthes (hookworm, strongyloidiasis, cutaneous larva migrans)). For food/waterborne diseases (e.g., bacterial diarrhea, hepatitis A, typhoid/paratyphoid fever, diarrhea-cholera, diarrhea-protozoal, brucellosis, hepatitis E), if ingesting local food and water, the health effects can temporarily incapacitate personnel (diarrhea) or result in prolonged illness (hepatitis A, typhoid/paratyphoid fever, brucellosis, hepatitis E). Risks from food/waterborne diseases may have been reduced with preventive medicine controls and mitigation, which includes hepatitis A and typhoid fever vaccinations and only drinking from approved water sources in accordance with standing CENTCOM policy. For other vector-borne endemic diseases (malaria, Crimean-Congo hemorrhagic fever, leishmaniasis-cutaneous (acute), sandfly fever, typhus-miteborne (scrub typhus)), these diseases may constitute a significant risk due to exposure to biting vectors; risk reduced to 'Low' by proper wear of the treated uniform, application of repellent to exposed skin, bed net use, and appropriate chemoprophylaxis, as well as minimizing areas of standing water and other vector-breeding areas. For water contact diseases (leptospirosis) activities involving extensive contact with surface water increase risk. For respiratory diseases (TB), personnel in close-quarter conditions could have been at risk for person-to-person spread. Animal contact diseases (rabies, Q fever), pose year-round risk. For soil transmitted helminthes (hookworm, strongyloidiasis, cutaneous larva migrans), risk may have been reduced by limiting exposure to soil contaminated with human or animal feces (including not sleeping on bare ground, and not walking barefoot).

Air quality: For inhalable coarse particulate matter less than 10 micrometers in diameter (PM₁₀) from environmental dust (including any burn pits and/or incinerators, which might have existed), the PM₁₀ overall short-term health risk was not evaluated due to no data for analysis. For inhalable fine particulate matter less than 2.5 micrometers in diameter (PM_{2.5}) from environmental dust (including any burn pits and/or incinerators, which might have existed), the PM_{2.5} overall short-term health risk was not evaluated due to no data for analysis. However, the Camp Panther and vicinity area is a dust-prone desert environment, with a semi-arid climate, also subject to vehicle traffic. Consequently, exposures to PM₁₀ and PM_{2.5} may vary, as conditions may vary, and may result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel while at this site, particularly exposures to high levels of dust such as during high winds or dust storms. For PM₁₀ and PM_{2.5}, certain subgroups of the deployed forces (e.g., those with pre-existing asthma/cardio-pulmonary conditions) are at greatest risk of developing notable health effects. Burn pits and/or incinerators might have existed in the vicinity (e.g., burn pits used by the local population); however, the PM₁₀ and the PM_{2.5} overall short-term health risks specifically for burn pits were not evaluated due to no environmental samples collected near burn pits provided for analysis— see Section 10.7. Where burn pits exist, exposures may vary, and exposures to high levels of PM₁₀ and PM_{2.5} from smoke may result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel and certain subgroups. Although most short-term health effects from exposure to particulate matter and incinerator and/or burn pit smoke should have resolved post-deployment, providers should be prepared to consider the relationship between deployment exposures and current complaints. Some individuals may have sought treatment for acute respiratory irritation while at Camp Panther and vicinity. Personnel who reported with symptoms or required treatment while at site(s) with burn pit activity should have exposure and treatment noted in medical record (e.g., electronic medical record and/or on a Standard Form (SF) 600 (*Chronological Record of Medical Care*)).

Long-term health risks & medical implications:

The following hazards may be associated with potential chronic health effects in some personnel during deployment at Camp Panther:

Air quality: For inhalable fine particulate matter less than 2.5 micrometers in diameter (PM_{2.5}) from environmental dust (including any burn pits and/or incinerators, which might have existed), the overall long-term health risk was not evaluated due to no data for analysis. Inhalable coarse particulate matter less than 10 micrometers in diameter (PM₁₀) from environmental dust (including any burn pits and/or incinerators, which might have existed) was not evaluated for long-term health risk due to no data for analysis and no available health guidelines. However, the Camp Panther and vicinity area is a dust-prone desert environment with a semi-arid climate, also subject to vehicle traffic, and conditions may have varied. Burn pits and/or incinerators might have existed in the vicinity (e.g., burn pits used by the local population); however, the PM₁₀ and the PM_{2.5} overall long-term health risks specifically for burn pits were not evaluated due to no environmental samples collected near burn pits provided for analysis— see Section 10.7. Where burn pits exist, exposures may vary, as conditions may have varied. For inhalational exposure to high levels of dust containing PM₁₀ and PM_{2.5}, such as during high winds or dust storms, and for exposures to incinerator and/or burn pit smoke, it is considered possible that some otherwise healthy personnel, who were exposed for a long-term period to dust and particulate matter, could develop certain health conditions

(e.g., reduced lung function, cardiopulmonary disease). Personnel with a history of asthma or cardiopulmonary disease could potentially be more likely to develop such chronic health conditions. While the dust and particulate matter exposures and exposures to burn pits are acknowledged, at this time there were no specific recommended, post-deployment medical surveillance evaluations or treatments. Providers should still consider overall individual health status (e.g., any underlying conditions/susceptibilities) and any potential unique individual exposures (such as burn pits/barrels, incinerators, occupational or specific personal dosimeter data) when assessing individual concerns. Certain individuals may need to be followed/evaluated for specific occupational exposures/injuries (e.g., annual audiograms as part of the medical surveillance for those enrolled in the Hearing Conservation Program; and personnel covered by Respiratory Protection Program and/or Hazardous Waste/Emergency Responders Medical Surveillance).

Table 2. Population-Based Health Risk Estimates – Camp Panther ^{1, 2}

Source of Identified Health Risk ³	Unmitigated Health Risk Estimate ⁴	Control Measures Implemented	Residual Health Risk Estimate ⁴
AIR			
Particulate matter less than 10 micrometers in diameter (PM ₁₀)	Short-term: No data were available for analysis to characterize health risk. Daily levels vary, acute health effects are (e.g., upper respiratory tract irritation) more pronounced during days with elevated PM levels. More serious effects are possible in susceptible persons (e.g., those with asthma/pre-existing respiratory diseases).	Limiting strenuous physical activities when air quality is especially poor; and actions such as closing tent flaps, windows, and doors.	Short-term: No data were available for analysis to characterize health risk. Daily levels vary, acute health effects (e.g., upper respiratory tract irritation) more pronounced during days with elevated PM levels. More serious effects are possible in susceptible persons (e.g., those with asthma/pre-existing respiratory diseases).
	Long-term: No health guidelines		Long-term: No health guidelines
Particulate matter less than 2.5 micrometers in diameter (PM _{2.5})	Short-term: No data were available for analysis to characterize health risk. Because Camp Panther is situated in a dusty, semi-arid desert environment, a majority of the time mild acute (short-term) health effects are anticipated. Elevated levels may produce mild eye, nose, or throat irritation in some personnel and pre-existing health conditions (e.g., asthma, or cardiopulmonary diseases) may be exacerbated.	Limiting strenuous physical activities when air quality is especially poor; taking actions such as closing tent flaps, windows, and doors.	Short-term: No data were available for analysis to characterize health risk. Because Camp Panther is situated in a dusty, semi-arid desert environment, a majority of the time mild acute (short-term) health effects are anticipated. Elevated levels may produce mild eye, nose, or throat irritation in some personnel and pre-existing health conditions (e.g., asthma, or cardiopulmonary diseases) may be exacerbated.
	Long-term: No data were available for analysis to characterize health risk. A small percentage of personnel may be at increased risk for developing chronic conditions, particularly those more susceptible to acute effects (e.g., those with asthma/pre-existing respiratory diseases).		Long-term: No data were available for analysis to characterize health risk. A small percentage of personnel may be at increased risk for developing chronic conditions, particularly those more susceptible to acute effects (e.g., those with asthma/pre-existing respiratory diseases).
ENDEMIC DISEASE			
Foodborne/ Waterborne (e.g., diarrhea- bacteriological)	Short-term: Variable; High (bacterial diarrhea, hepatitis A, typhoid fever) to Moderate (diarrhea-cholera, diarrhea- protozoal, brucellosis, hepatitis E) to Low (polio) if ingesting local food/water, the health effects can temporarily incapacitate personnel (diarrhea) or result in prolonged illness (hepatitis A, typhoid fever, hepatitis E, brucellosis).	Preventive measures include hepatitis A and typhoid fever vaccination and consumption of food and water only from approved sources.	Short-term: Low to none
	Long-term: None identified		Long-term: No data available
Arthropod Vector- Borne	Short-term: Variable; High for malaria and Crimean-Congo hemorrhagic fever, Moderate for leishmaniasis-cutaneous (acute), sandfly fever, scrub typhus-miteborne; and Low for the plague and West Nile fever.	Preventive measures include proper wear of treated uniform, application of repellent to exposed skin, bed net use, minimizing areas of standing water and appropriate chemoprophylaxis.	Short-term: Low
	Long-term: Low for Leishmaniasis-visceral infection.		Long-term: No data available
Water-Contact	Short-term: Moderate for leptospirosis	Recreational swimming in	Short-term: Low for leptospirosis.

Source of Identified Health Risk ³	Unmitigated Health Risk Estimate ⁴	Control Measures Implemented	Residual Health Risk Estimate ⁴
(e.g., wading, swimming)	Long-term: No data available	surface waters not likely in this area of Afghanistan during this time period.	Long-term: No data available
Respiratory	Short-term: Variable; Moderate for tuberculosis (TB) to Low for meningococcal meningitis and Middle East respiratory syndrome coronavirus (MERS-CoV).	Providing adequate living and work space; medical screening.	Short-term: Low
	Long-term: No data available		Long-term: No data available
Animal Contact	Short-term: Variable; Moderate for rabies, Q-fever, to Low for anthrax, and avian influenza.	Prohibiting contact with, adoption, or feeding of feral animals IAW U.S. Central Command (CENTCOM) General Order (GO) 1C. Risks are further reduced in the event of assessed contact by prompt post-exposure rabies prophylaxis IAW The Center for Disease Control's (CDC) Advisory Committee on Immunization Practices guidance.	Short-term: No data available
	Long-term: Low (Rabies)		Long-term: No data available
Soil-transmitted	Short-term: Moderate for soil transmitted helminthes (hookworm, strongyloidiasis, cutaneous larva migrans).	Risk was reduced to Low by limiting exposure to soil contaminated with human or animal feces (including sleeping on bare ground and walking barefoot).	Short-term: Low
	Long-term: No data available		Long-term: No data available
VENOMOUS ANIMAL/ INSECTS			
Snakes, scorpions, and spiders	Short-term: Low; If encountered, effects of venom vary with species from mild localized swelling (e.g., <i>Latrodectus dahlia</i>) to potentially lethal effects (e.g. <i>Gloydius halys</i>).	Risk reduced by avoiding contact, proper wear of uniform (especially footwear), and proper and timely treatment.	Short-term: Low; If encountered, effects of venom vary with species from mild localized swelling (e.g., <i>Latrodectus dahlia</i>) to potentially lethal effects (e.g. <i>Gloydius halys</i>).
	Long-term: No data available		Long-term: No data available
HEAT/COLD STRESS			
Heat	Short-term: Not evaluated	Work-rest cycles, proper hydration and nutrition, and Wet Bulb Globe Temperature (WBGT) monitoring.	Short-term: Not evaluated
	Long-term: Low; However, the risk may be greater to certain susceptible persons—those older (i.e., greater than 45 years), in lesser physical shape, or with underlying medical/health conditions.		Long-term: Low; However, the risk may be greater to certain susceptible persons—those older (i.e., greater than 45 years), in lesser physical shape, or with underlying medical/health conditions.
Cold	Short-term: Not evaluated.	Risks from cold stress reduced with protective measures such as use of the buddy system, limiting exposure during cold weather, proper hydration and nutrition, and proper wear of issued protective clothing.	Short-term: Not evaluated.
	Long-term: Low; Long-term health implications from cold injuries are rare but can occur, especially from more serious injuries such as frost bite.		Long-term: Low; Long-term health implications from cold injuries are rare but can occur, especially from more serious injuries such as frost bite.

Source of Identified Health Risk ³	Unmitigated Health Risk Estimate ⁴	Control Measures Implemented	Residual Health Risk Estimate ⁴
Unique Incidents/ Concerns			
Burn Pits	Short-term: Burn pits and/or incinerators might have existed at Camp Panther (for example, burn pits used by the local population); however, there are no reports or sampling data to indicate their presence or absence. Consequently, the PM ₁₀ and the PM _{2.5} overall short-term health risks specifically for burn pits were not evaluated – see Section 10.7. A majority of the time mild acute (short term) health effects are anticipated; certain peak levels may produce mild eye, nose, or throat irritation in some personnel and pre-existing health conditions (e.g., asthma, or cardiopulmonary diseases) may be exacerbated.	Risks reduced by limiting strenuous physical activities when air quality was especially poor; and action such as closing tent flaps, windows, and doors. Other control measures included locating burn pits downwind of camps, increased distance from troop populations, and improved waste segregation and management techniques.	Short-term: Burn pits and/or incinerators might have existed at Camp Panther (for example, burn pits used by the local population); however, there are no reports or sampling data to indicate their presence or absence. Consequently, the PM ₁₀ and the PM _{2.5} overall short-term health risks specifically for burn pits were not evaluated – see Section 10.7. A majority of the time mild acute (short term) health effects are anticipated; certain peak levels may produce mild eye, nose, or throat irritation in some personnel and pre-existing health conditions (e.g., asthma, or cardiopulmonary diseases) may be exacerbated.
	Long-term: Burn pits and/or incinerators might have existed at Camp Panther (for example, burn pits used by the local population); however, there are no reports or sampling data to indicate their presence or absence. Consequently, the PM ₁₀ and the PM _{2.5} overall long-term health risks specifically for burn pits were not evaluated – see Section 10.7. Exposure to burn pit smoke is variable. Exposure to high levels of PM ₁₀ and PM _{2.5} in the smoke may be associated with some otherwise healthy personnel, who were exposed for a long-term period, possibly developing certain health conditions (e.g., reduced lung function, cardiopulmonary disease). Personnel with a history of asthma or cardiopulmonary disease could potentially be more likely to develop such chronic health conditions.		Long-term: Burn pits and/or incinerators might have existed at Camp Panther (for example, burn pits used by the local population); however, there are no reports or sampling data to indicate their presence or absence. Consequently, the PM ₁₀ and the PM _{2.5} overall long-term health risks specifically for burn pits were not evaluated – see Section 10.7. Exposure to burn pit smoke is variable. Exposure to high levels of PM ₁₀ and PM _{2.5} in the smoke may be associated with some otherwise healthy personnel, who were exposed for a long-term period, possibly developing certain health conditions (e.g., reduced lung function, cardiopulmonary disease). Personnel with a history of asthma or cardiopulmonary disease could potentially be more likely to develop such chronic health conditions.

¹This Summary Table provides a qualitative estimate of population-based short- and long-term health risks associated with the occupational environment conditions at Camp Panther or Afghanistan. It does not represent an individual exposure profile. Actual individual exposures and health effects depend on many variables. For example, while a chemical may have been present in the environment, if a person did not inhale, ingest, or contact a specific dose of the chemical for adequate duration and frequency, then there may have been no health risk. Alternatively, a person at a specific location may have experienced a unique exposure which could result in a significant individual exposure. Any such person seeking medical care should have their specific exposure documented in an SF600.

²This assessment is based on specific environmental sampling data and reports obtained in 2008. Sampling locations are assumed to be representative of exposure points for the camp population but may not reflect all the fluctuations in environmental quality or capture unique exposure incidents.

³This Summary Table is organized by major categories of identified sources of health risk. It only lists those sub-categories specifically identified and addressed at Camp Panther or Afghanistan when site specific information is not available. The health risks are presented as Low, Moderate, High or Extremely High for both acute and chronic health effects. The health risk level is based on an assessment of both the potential severity of the health effects that could be caused and probability of the exposure that would produce such health effects. Details can be obtained from the Army Public Health Center (APHC). Where applicable, "None Identified" is used when though a potential exposure is identified, and no health risks of either a specific acute or chronic health effects are determined. More detailed descriptions of OEH exposures that are evaluated but determined to pose no health risk are discussed in the following sections of this report.

⁴Health risks in this Summary Table are based on quantitative surveillance thresholds (e.g., endemic disease rates; host/vector/pathogen surveillance) or screening levels, e.g., Military Exposure Guidelines (MEGs) for chemicals. Some previous assessment reports may provide slightly inconsistent health risk estimates because quantitative criteria such as MEGs may have changed since the samples were originally evaluated and/or because this assessment makes use of all historic site data while previous reports may have only been based on a select few samples.

1 Discussion of Health Risks at Camp Panther, Afghanistan by Source

The following sections provide additional information about the OEH conditions summarized above. All risk assessments were performed using the methodology described in the U.S. Army Public Health Center (USAPHC) Technical Guide 230, *Environmental Health Risk Assessment and Chemical Exposure Guidelines for Deployed Military Personnel* (Reference 4). All OEH risk estimates represent residual risk after accounting for preventive controls in place. Occupational exposures and exposures to endemic diseases are greatly reduced by preventive measures. For environmental exposures related to airborne dust, there are limited preventive measures available, and available measures have little efficacy in reducing exposure to ambient conditions.

2 Air

2.1 Site-Specific Sources Identified

Although the exact location of Camp Panther is unknown, Afghanistan is a dusty, semi-arid desert environment. Inhalational exposure to high levels of dust and particulate matter, such as during high winds or dust storms, may result in mild to more serious short-term health effects (e.g., eye, nose or throat and lung irritation) in some personnel. Additionally, certain subgroups of the deployed forces (e.g., those with pre-existing asthma/cardio pulmonary conditions) are at greatest risk of developing notable health effects.

2.2 Particulate matter

Particulate matter (PM) is a complex mixture of extremely small particles suspended in the air. The PM includes solid particles and liquid droplets emitted directly into the air by sources such as: power plants, motor vehicles, aircraft, generators, construction activities, fires, and natural windblown dust. The PM can include sand, soil, metals, volatile organic compounds (VOC), allergens, and other compounds such as nitrates or sulfates that are formed by condensation or transformation of combustion exhaust. The PM composition and particle size vary considerably depending on the source. Generally, PM of health concern is divided into two fractions: PM₁₀, which includes coarse particles with a diameter of 10 micrometers or less, and fine particles less than 2.5 micrometers (PM_{2.5}), which can reach the deepest regions of the lungs when inhaled. Exposure to excessive PM is linked to a variety of potential health effects.

2.3 Particulate matter, less than 10 micrometers (PM₁₀)

2.3.1 Exposure Guidelines:

Short Term (24-hour) PM₁₀ (micrograms per cubic meter, µg/m³):

- Negligible MEG = 250
- Marginal MEG = 420
- Critical MEG = 600

Long-term PM₁₀ MEG (µg/m³):

- Not defined and not available.

2.3.2 Sample data/Notes:

No valid PM₁₀ air samples were available for analysis.

2.3.3 Short-term health risks:

Not evaluated.

2.3.4 Long-term health risk:

Not Evaluated-no data nor available health guidelines. The U.S. Environmental Protection Agency (EPA) has retracted its long-term standard (National Ambient Air Quality Standards, NAAQS) for PM₁₀ due to an inability to clearly link chronic health effects with chronic PM₁₀ exposure levels.

2.4 Particulate Matter, less than 2.5 micrometers (PM_{2.5})

2.4.1 Exposure Guidelines:

Short-Term (24-hour) PM_{2.5} (µg/m³):

- Negligible MEG = 65
- Marginal MEG = 250
- Critical MEG = 500

Long-term (1 year) PM_{2.5} MEGs (µg/m³):

- Negligible MEG = 15
- Marginal MEG = 65.

2.4.2 Sample data/Notes:

No valid PM_{2.5} air samples were available for analysis.

2.4.3 Short- and Long-term health risks:

Not evaluated.

2.5 Airborne Metals

2.5.1 Sample data/Notes:

No valid airborne metal samples were available for analysis.

2.5.2 Short- and Long-term health risks:

Not evaluated.

2.6 Volatile Organic Compounds (VOC)

2.6.1 Sample data/Notes:

No valid VOC samples were available for analysis.

2.6.2 Short- and long-term health risks:

Not evaluated.

3 Soil

3.1 Site-Specific Sources Identified

3.2 Sample data/Notes:

Three valid discreet surface soil samples were collected from an area referred to as “open grounds” on 30 August 2008 to assess OEH health risk to deployed personnel. The primary soil contamination exposure pathways are dermal contact and dust inhalation. Typical parameters analyzed for included: semi volatile organic compounds (SVOCs), heavy metals, polychlorinated biphenyls (PCBs), pesticides, and herbicides. For the risk assessment, personnel are assumed to remain at this location for greater than 1 year.

3.3 Short-term health risk:

Not an identified source of health risk. Currently, sampling data for soil are not evaluated for short-term (acute) health risks.

3.4 Long-term health risk:

None identified based on available sample data. No parameters exceeded its 1-year Negligible MEGs.

4 Water

No reverse osmosis water purification unit (ROWPU) treated, disinfected fresh bulk water, bottled water, or untreated well water samples were available for analysis.

4.1 Short and long-term health risk

Not evaluated

5 Military Unique

5.1 Chemical Biological, Radiological Nuclear (CBRN) Weapons

No specific hazard sources were documented in the Defense Occupational and Environmental Health Readiness System (DOEHRS) in 2008 (Reference 1).

5.2 Depleted Uranium (DU)

No specific hazard sources were documented in the DOEHRS in 2008 (Reference 1).

5.3 Ionizing Radiation

No specific hazard sources were documented in the DOEHRS in 2008 (Reference 1).

5.4 Non-Ionizing Radiation

No specific hazard sources were documented in the DOEHRS in 2008 (Reference 1).

6 Endemic Diseases

This document lists the endemic diseases reported in the region, its specific health risks and severity and general health information about the diseases. CENTCOM Modification (MOD) 9 (Reference 6) lists deployment requirements, to include immunizations and chemoprophylaxis, in effect during the timeframe of this POEMS.

6.1 Foodborne and Waterborne Diseases

Foodborne and waterborne diseases in the area are transmitted through the consumption of local food and water. Local unapproved food and water sources (including ice) are heavily contaminated with pathogenic bacteria, parasites, and viruses to which most U.S. Service Members have little or no natural immunity. Effective disease surveillance has been improved to cover the majority of the country since 2009. There is still underreporting of specific disease incidence. Diarrheal diseases are expected to temporarily incapacitate a very high percentage of U.S. personnel within days if local food, water, or ice is consumed. Hepatitis A and typhoid fever infections typically cause prolonged illness in a smaller percentage of unvaccinated personnel. Vaccinations are required for DoD personnel and contractors. In addition, although not specifically assessed in this document, significant outbreaks of viral gastroenteritis (e.g., norovirus) and food poisoning (e.g., *Bacillus cereus*, *Clostridium perfringens*, *Staphylococcus* spp.) may occur. Key disease risks are summarized below:

Mitigation strategies were in place and included consuming food and water from approved sources, vaccinations (when available), frequent hand washing and general sanitation practices.

6.1.1 Diarrheal diseases (bacteriological)

High, mitigated to Low: Diarrheal diseases are expected to temporarily incapacitate a very high percentage of personnel (potentially over 50% per month) within days if local food, water, or ice is consumed. Field conditions (including lack of hand washing and primitive sanitation) may facilitate person-to-person spread and epidemics. Typically mild disease treated in outpatient setting; recovery and return to duty in less than 72 hours with appropriate therapy. A small proportion of infections may require greater than 72 hours limited duty or hospitalization.

6.1.2 Hepatitis A, typhoid/paratyphoid fever, and diarrhea-protozoal

High, mitigated to Low: Unmitigated health risk to U.S. personnel is high year round for hepatitis A and typhoid/paratyphoid fever, and Moderate for diarrhea-protozoal. Mitigation was in place to reduce the risks to Low. Hepatitis A, typhoid/paratyphoid fever, and diarrhea-protozoal disease may cause prolonged illness in a small percentage of personnel (less than 1% per month). Although much rarer, other potential diseases in this area that are also considered a Moderate risk include: hepatitis E, diarrhea-cholera, and brucellosis.

6.1.3 Polio

Low: Potential health risk to U.S. personnel is Low. Despite a concerted global eradication campaign, poliovirus continues to affect children and adults in Afghanistan. Polio is a highly infectious disease that invades the nervous system. The virus is transmitted by person-to-person, typically by hands, food or water contaminated with fecal matter or through direct contact with the infected person's saliva. An infected person may spread the virus to others immediately before and about 1 to 2 weeks after symptoms appear. The virus can live in an infected person's feces for many weeks. About 90% of people infected have no symptoms, and about 1% have a very severe illness leading to muscle weakness, difficulty breathing, paralysis, and sometimes death. People who do not have symptoms can still pass the virus to others and make them sick.

6.1.4 Short-term Health Risks:

Low: The overall unmitigated short-term risk associated with foodborne and waterborne diseases are considered High (bacterial diarrhea, hepatitis A, typhoid/paratyphoid fever) to Moderate (diarrhea-cholera, diarrhea-protozoal, brucellosis, hepatitis E) to Low (polio) if local food or water is consumed. Preventive Medicine measures reduced the risk to Low. Confidence in the health risk estimate is medium.

6.1.5 Long-term Health Risks:

None identified based on available data.

6.2 Arthropod Vector-Borne Diseases

During the warmer months, the climate and ecological habitat support populations of arthropod vectors, including mosquitoes, ticks, mites, and sandflies. Significant disease transmission is sustained countrywide, including urban areas. Malaria, the major vector-borne health risk in Afghanistan, is capable of debilitating a high percentage of personnel for up to a week or more. Mitigation strategies were in place and included proper wear of treated uniforms, application of repellent to exposed skin, and use of bed nets and chemoprophylaxis (when applicable). Additional methods included the use of pesticides, reduction of pest/breeding habitats, and engineering controls.

6.2.1 Malaria

High, mitigated to Low: Potential unmitigated risk to U.S. personnel is High during warmer months (typically April through November) but reduced to Low with mitigation measures. Malaria incidents are often associated with the presence of agriculture activity, including irrigation systems and standing water, which provide breeding habitats for vectors. A small number of cases may occur among personnel exposed to mosquito (*Anopheles* spp.) bites. Malaria incidents may cause debilitating febrile illness typically requiring 1 to 7 days of inpatient care, followed by return to duty. Severe cases may require intensive care or prolonged convalescence.

6.2.2 Leishmaniasis

Moderate, mitigated to Low: The disease risk is Moderate during the warmer months when sandflies are most prevalent, but reduced to low with mitigation measures. Leishmaniasis is transmitted by sand flies. A small number of cases (less than 1% per month attack rate) could occur among personnel exposed to sandfly bites in areas with infected people, rodents, dogs, or other reservoir animals. In groups of personnel exposed to heavily infected sandflies in focal areas, attack rates can be very high (over 50%). There are two forms of the disease; cutaneous (acute form) and visceral (a more latent form of the disease). The leishmaniasis parasites may survive for years in infected individuals and this infection may go unrecognized by physicians in the U.S. when infections become symptomatic years later. Cutaneous infection is unlikely to be debilitating, though lesions may be disfiguring. Visceral leishmaniasis disease can cause severe febrile illness which typically requires hospitalization with convalescence over 7 days.

6.2.3 Crimean-Congo hemorrhagic fever

High, mitigated to Low: Unmitigated risk is High, but reduced to low with mitigation measures. Crimean-Congo hemorrhagic fever occurs in a small number of cases (less than 1% per month attack rate) and is transmitted by tick bites or occupational contact with blood or secretions from infected animals. The disease typically requires intensive care with fatality rates from 5% to 50%.

6.2.4 Sandfly fever

Moderate, mitigated to Low: Sandfly fever has a Moderate risk with potential disease rates from 1% to 10% per month; under worst case conditions disease rates can be as high as 50%. Mitigation measures reduced the risk to low. The disease is transmitted by sandflies and occurs more commonly in children though adults are still at risk. Sandfly fever disease typically resulted in debilitating febrile illness requiring 1 to 7 days of supportive care followed by return to duty.

6.2.5 Plague

Low: Potential health risk to U.S. personnel is Low year round. Bubonic plague typically occurred as sporadic cases among people who come in contact with wild rodents and their fleas during work, hunting, or camping activities. Outbreaks of human plague are rare and typically occur in crowded urban settings associated with large increases in infected commensal rats (*Rattus rattus*) and their flea populations. Some untreated cases of bubonic plague may develop into secondary pneumonic plague. Respiratory transmission of pneumonic plague is rare but has the potential to cause significant outbreaks. Close contact is usually required for transmission. In situations where respiratory transmission of plague is suspected, weaponized agent must be considered. Extremely rare cases (less than 0.01% per month attack rate) could occur. Incidence could result in potentially severe illness which may require more than 7 days of hospitalization and convalescence.

6.2.6 Typhus-miteborne (scrub typhus)

Moderate, mitigated to Low: Potential health risk to U.S. personnel is Moderate during warmer months (typically March through November) when vector activity is highest. Mitigation measures reduced the risk to low. Mite-borne typhus is a significant cause of febrile illness in local populations with rural exposures in areas where the disease is endemic. Large outbreaks have occurred when non-indigenous personnel such as military forces enter areas with established local transmission. The disease is transmitted by the larval stage of trombiculid mites (chiggers), which are typically found in areas of grassy or scrubby vegetation, often in areas which have undergone clearing and regrowth. Habitats may include sandy beaches, mountain deserts, cultivated rice fields, and rain forests. Although data are insufficient to assess potential disease rates, attack rates can be very high (over 50%) in groups of personnel exposed to heavily infected "mite islands" in focal areas. The disease can cause debilitating febrile illness typically requiring 1 to 7 days of inpatient care, followed by return to duty.

6.2.7 West Nile fever

Low: West Nile fever is present. The disease is maintained by the bird population and transmitted to humans via mosquito vector. Typically, infections in young, healthy adults were asymptomatic although fever, headache, tiredness, body aches (occasionally with a skin rash on trunk of body), and swollen lymph glands can occur. This disease is associated with a low risk estimate.

6.2.9 Short-term health risks:

Low: The unmitigated health risk estimate is High for malaria and Crimean-Congo hemorrhagic fever (infection rate of less than 1% per month), Moderate for leishmaniasis-cutaneous (acute), sandfly fever, typhus-miteborne; and Low for, the plague and West Nile fever. Health risk is reduced to low by proper wear of the uniform, application of repellent to exposed skin, and appropriate chemoprophylaxis. Confidence in health risk estimate was medium.

6.2.10 Long-term health risks:

Moderate, mitigated to Low: The unmitigated risk is moderate for leishmaniasis-visceral (chronic). Risk is reduced to Low by proper wear of the uniform and application of repellent to exposed skin. Confidence in the risk estimate is medium.

6.3 Water Contact Diseases

Operations or activities that involve extensive water contact may result in personnel being temporarily debilitated with leptospirosis in some locations. Leptospirosis health risk typically increases during flooding. In addition, although not specifically assessed in this document, bodies of surface water are likely to be contaminated with human and animal waste. Activities such as wading or swimming may result in exposures to enteric diseases such as diarrhea and hepatitis via incidental ingestion of water. Prolonged water contact also may lead to the development of a variety of potentially debilitating skin conditions such as bacterial or fungal dermatitis. Mitigation strategies were in place and included avoiding water contact and recreational water activities, proper wear of uniform (especially footwear), and protective coverings for cuts/abraded skin.

6.3.1 Leptospirosis

Moderate, mitigated to Low: Human infections occur seasonally (typically April through November) through exposure to water or soil contaminated by infected animals and is associated with wading, and swimming in contaminated, untreated open water. The occurrence of flooding after heavy rainfall facilitates the spread of the organism because as water saturates the environment, *Leptospira* spp. present in the soil passes directly into surface waters. *Leptospira* spp. can enter the body through cut or abraded skin, mucous membranes, and conjunctivae. Infection may also occur from ingestion of contaminated water. The acute, generalized illness associated with infection may mimic other tropical diseases (for example, dengue fever, malaria, and typhus), and common symptoms include fever, chills, myalgia, nausea, diarrhea, cough, and conjunctival suffusion. Manifestations of severe disease can include jaundice, renal failure, hemorrhage, pneumonitis, and hemodynamic collapse. Recreational activities involving extensive water contact may result in personnel being temporarily debilitated with leptospirosis. Incidence could result in debilitating febrile illness typically requiring 1 to 7 days of inpatient care, followed by return to duty; some cases may require prolonged convalescence. This disease is associated with a Moderate health risk estimate.

6.3.2 Short-term health risks:

Low: Unmitigated Health risk of leptospirosis is Moderate during warmer months. Mitigation measures reduce the risk to Low. Confidence in the health risk estimate is medium.

6.3.3 Long-term health risks:

None identified based on available data.

6.4 Respiratory Diseases

Although not specifically assessed in this document, deployed U.S. Forces may be exposed to a wide variety of common respiratory infections in the local population. These include influenza, pertussis, viral upper respiratory infections, viral and bacterial pneumonia, measles, and others. The U.S. military populations living in close-quarter conditions are at risk for substantial person-to-person spread of respiratory pathogens. Influenza is of particular concern because of its ability to debilitate large numbers of unvaccinated personnel for several days. Mitigation strategies were in place and included

routine medical screenings, vaccination, enforcing minimum space allocation in housing units, implementing head-to-toe sleeping in crowded housing units, implementation of proper personal protective equipment (PPE) when necessary for healthcare providers and detention facility personnel.

6.4.1 Tuberculosis (TB)

Moderate, mitigated to Low: Potential health risk to U.S. personnel is Moderate, mitigated to Low, year round. Transmission typically requires close and prolonged contact with an active case of pulmonary or laryngeal TB, although it also can occur with more incidental contact. Individuals with prolonged indoor exposure to the local population are at increased risk for latent TB infection.

6.4.2 Meningococcal meningitis

Low: Meningococcal meningitis poses a Low risk and is transmitted from person to person through droplets of respiratory or throat secretions. Close and prolonged contact facilitates the spread of this disease. Meningococcal meningitis is potentially a very severe disease typically requiring intensive care; fatalities may occur in 5-15% of cases.

6.4.3 Middle East respiratory syndrome coronavirus (MERS-CoV)

Low: Although no cases have been reported in Afghanistan, Middle East respiratory syndrome coronavirus (MERS-CoV) is known to occur within the region. Most MERS patients developed severe acute respiratory illness with symptoms of fever, cough and shortness of breath. MERS-CoV has spread from ill people to others through close contact, such as caring for or living with an infected person. The incubation period for MERS-CoV is usually about 5 to 6 days, but can range from 2 to 14 days. Currently, there is no vaccine to prevent MERS-CoV infection.

6.4.4 Short-term health risks:

Low: Moderate (TB) to Low (for meningococcal meningitis and MERS-CoV). Overall risk was reduced to Low with mitigation measures. Confidence in the health risk estimate is medium.

6.4.5 Long-term health risks:

None identified based on available data. Tuberculosis is evaluated as part of the post deployment health assessment (PDHA). A TB skin test is required post-deployment if potentially exposed and is based upon individual service policies.

6.5 Animal-Contact Diseases

6.5.1 Rabies

Moderate, mitigated to Low: Rabies posed a year-round moderate risk. Occurrence in local animals was well above U.S. levels due to the lack of organized control programs. Dogs are the primary reservoir of rabies in Afghanistan, and a frequent source of human exposure. Rabies is transmitted by exposure to the virus-laden saliva of an infected animal, typically through bites, but could occur from scratches contaminated with the saliva. A U.S. Army Soldier deployed to Afghanistan from May 2010 to May 2011 died of rabies in New York on 31 August 2011 (Reference 7). Laboratory results indicated the Soldier was infected from contact with a dog while deployed. Although the vast majority (>99%) of persons who develop rabies disease will do so within a year after a risk exposure, there have been rare reports of individuals presenting with rabies disease up to six years or more after their last known risk exposure. Mitigation strategies included command emphasis of CENTCOM GO 1C, reduction of

animal habitats, active pest management programs, and timely treatment of feral animal scratches/bites.

6.5.2 Anthrax

Low: Anthrax cases are rare in indigenous personnel, and pose a Low risk to U.S. personnel. Anthrax is a naturally occurring infection; cutaneous anthrax is transmitted by direct contact with infected animals or carcasses, including hides. Eating undercooked infected meat may result in contracting gastrointestinal anthrax. Pulmonary anthrax is contracted through inhalation of spores and is extremely rare. Mitigation measures included consuming approved food sources, proper food preparation and cooking temperatures, avoidance of animals and farms, dust abatement when working in these areas, vaccinations, and proper PPE for personnel working with animals.

6.5.3 Q-Fever

Moderate, mitigated to Low: Potential health risk to U.S. personnel is Moderate, but mitigated to Low, year round. Rare cases are possible among personnel exposed to aerosols from infected animals, with clusters of cases possible in some situations. Significant outbreaks (affecting 1-50%) can occur in personnel with heavy exposure to barnyards or other areas where animals are kept. Unpasteurized milk may also transmit infection. The primary route of exposure is respiratory, with an infectious dose as low as a single organism. Incidence could result in debilitating febrile illness, sometimes presenting as pneumonia, typically requiring 1 to 7 days of inpatient care followed by return to duty. Mitigation strategies in place as listed in paragraph 6.5.2 except for vaccinations.

6.5.4 Avian influenza

Low: Potential health risk to U.S. personnel is Low. Although avian influenza is easily transmitted among birds, bird-to-human transmission is extremely inefficient. Human-to-human transmission appears to be exceedingly rare, even with relatively close contact. Extremely rare cases (less than 0.01% per month attack rate) could occur. Incidence could result in very severe illness with fatality rate higher than 50 percent in symptomatic cases. Mitigation strategies included avoidance of birds/poultry and proper cooking temperatures for poultry products.

6.5.5 Short-term health risks:

Low: The short-term unmitigated risk is Moderate for rabies and Q-fever, to Low for anthrax and avian influenza. Mitigation measures reduced the overall risk to Low. Confidence in risk estimate is medium.

6.5.6 Long-term health risks:

Low: A Low long term risk exists for rabies because in rare cases, the incubation period for rabies can be several years.

6.6 Soil-transmitted helminths (hookworm, strongyloidiasis, cutaneous larva migrans)

Moderate, mitigated to Low: Potential health risk to U.S. personnel is Moderate during warmer months (typically March through November) when vector activity is highest. Mitigation measures reduced the risk to low. A small number of cases (less than 1% per month attack rate) could occur among personnel with direct skin exposure to soil contaminated with human or animal feces (including sleeping on bare ground and walking barefoot). Initial skin symptoms typically are mild and are not debilitating. However, systemic symptoms of fever, cough, abdominal pain, nausea, and diarrhea may develop weeks to months after initial infection with hookworm or *Strongyloides* spp. More severe

infections with high worm burden may be debilitating in some cases. Rates of infection in U.S. personnel will be highly variable, depending on specific local environmental conditions. Rates of infection in U.S. personnel are expected to be less than 1 percent per month in most locations. However, rates in some focal areas with heavily contaminated soil could exceed 1 percent per month.

6.6.1 Short-term health risks:

Low: Moderate for soil transmitted helminthes. Overall risk was reduced to Low with mitigation measures. Confidence in the health risk estimate is medium.

6.6.2 Long-term health risks:

None identified based on available data.

7 Venomous Animals

All information was taken directly from the Armed Forces Pest Management Board (Reference 8) and the Clinical Toxinology Resources web site from the University of Adelaide, Australia (Reference 9). The precise location of Camp Panther was not provided in the DOEHRS record. The species listed below have home ranges in Afghanistan and may be found in the location of Camp Panther and potentially present a health risk if they are encountered by personnel. See Section 10 for more information about pesticides and pest control measures.

7.1 Spiders

- *Latrodectus dahlia* (widow spider): Severe envenoming possible, potentially lethal. However, venom effects are mostly minor and even significant envenoming is unlikely to be lethal.

7.2 Scorpions

- *Androctonus afghanus*, *Androctonus amoreuxi* and *Androctonus baluchicus*: Severe envenoming possible, potentially lethal. Severe envenoming may produce direct or indirect cardio toxicity, with cardiac arrhythmias and cardiac failure. Hypovolaemic hypotension possible in severe cases due to fluid loss through vomiting and sweating.

- *Afghanobuthus nuamanni*, *Buthacus striffleri*, *Compsobuthus afghanus*, *Compsobuthus rugosulus*, *Compsobuthus tofti*, *Mesobuthus caucasicus*, *Mesobuthus eupeus*, *Mesobuthus macmahoni*, *Orthochirus afghanus*, *Orthochirus bicolor*, *Orthochirus danielleae*, *Orthochirus erardi*, *Orthochirus heratensis*, *Orthochirus jalalabadensis*, *Orthochirus monodi*, *Orthochirus pallidus*, *Orthochirus samrchelsis*, *Orthochirus scrobiculosus*, and *Sassanidotus gracilis*: There are a number of dangerous Buthid scorpions, but there are also some known to cause minimal effects only. Without clinical data it is unclear where these species fit within that spectrum.

- *Hottentotta alticola* and *Hottentotta saulcyi*: Moderate envenoming possible but unlikely to prove lethal. Stings by these scorpions are likely to cause only short lived local effects, such as pain, without systemic effects.

- *Scorpiops afghanus*, *Scorpiops lindbergi*: Mild envenoming only, not likely to prove lethal. Stings by these scorpions are likely to cause only short lived local effects, such as pain, without systemic effects.

7.3 Snakes

- *Echis carinatus multisquamatus* (central Asian saw-scaled viper), *Echis carinatus sochureki* (Sochurek's saw-scaled viper), *Gloydius halys* (Haly's pit viper): Severe envenoming possible, potentially lethal. Bites may cause moderate to severe coagulopathy and haemorrhagins causing extensive bleeding.
- *Eristocophis mcmahoni* (McMahon's viper): Severe envenoming possible, potentially lethal. Venom shows strong hemorrhagic activity. Mild to Moderate neurotoxic effects may occur.
- *Macrovipera lebetina obtuse* (Levantine viper), and *Macrovipera lebetina turanica* (Levantine viper): Severe envenoming possible, potentially lethal. Bites may cause mild to severe local effects, shock & coagulopathy.
- *Naja oxiana* (oxus cobra): Severe envenoming possible, potentially lethal. Bites can cause systemic effects, principally flaccid paralysis.
- *Pseudocerastes persicus* (Persian horned viper): Unlikely to cause significant envenoming; limited clinical data suggest bites result in local effects only.
- *Bungarus caeruleus* (Common krait): Severe envenoming likely, high lethality potential. Krait bites can cause moderate to severe flaccid paralysis, respiratory failure, requiring intubation & ventilation in severe cases. Most victims bitten while asleep in huts at night. Bites may produce invisible or barely perceptible puncture marks. Human mortality rate is high without use of antivenom. Antivenom may prevent worsening of paralysis, but may not reverse established paralysis.
- *Gloydius himalayanus* (Himalayan pit viper), *Gloydius intermedius* (Central Asian pit viper): Potentially lethal envenoming, though unlikely, cannot be excluded. Bites cause in local and sometimes systemic effects including necrosis, coagulopathy, and renal failure.

7.4 Short-term health risk:

Low: If encountered, effects of venom vary with species from mild localized swelling (e.g. *Latrodectus dahlia*) to potentially lethal effects (e.g., *Gloydius halys*). See effects of venom above. Mitigation strategies included avoiding contact, proper wear of uniform (especially footwear), and timely medical treatment. Confidence in the health risk estimate is low (Reference 5, Table 3-6).

7.5 Long-term health risk:

None identified based on available data.

8 Heat/Cold Stress

8.1 Heat

The precise location of Camp Panther is not known. Therefore historical climate data was not available. As with other locations in Afghanistan, it is expected that summer months are hottest and tend to be dry. Additionally, work intensity and clothing/equipment worn posed a greater health risk of heat stress injury than environmental factors alone (Reference 10). Managing risk of hot weather operations included monitoring work/rest periods, proper hydration, and taking individual risk factors (e.g., acclimation, weight, and physical condition) into consideration. Risk of heat stress/injury was

reduced with preventive measures.

8.1.1 Short-term health risk:

Not evaluated: Based on standard Army policy the risk of heat injury should have been reduced to low through preventive measures such as work/rest cycles, proper hydration and nutrition, and monitoring Wet Bulb Globe Temperature (WBGT). Risk of heat injury in unacclimatized or susceptible populations (older, previous history of heat injury, poor physical condition, underlying medical/health conditions), and those under operational constraints (equipment, PPE, vehicles) may have higher risk associated with heat exposure.

8.1.2 Long-term health risk:

Low: The long-term risk is Low. However, the risk may be greater for certain susceptible persons—those older (i.e., greater than 45 years), in lesser physical shape, or with underlying medical/health conditions. Long-term health implications from heat injuries are rare but may occur, especially from more serious injuries such as heat stroke. It is possible that high heat in conjunction with various chemical exposures may increase long-term health risks, though specific scientific evidence is not conclusive. Confidence in these risk estimates is medium (Reference 5, Table 3-6).

8.2 Cold

The precise location of Camp Panther is not known. Therefore historical climate data was not available. As with other locations in Afghanistan it is expected that winter months are the coldest and tend to receive the most precipitation. However, personnel may encounter significantly lower temperatures during field operations at higher altitudes. As with heat stress/injuries, cold stress/injuries are largely dependent on operational and individual factors instead of environmental factors alone (Reference 10).

8.2.1 Short term health risk:

Not Evaluated

8.1.2 Long-term health risk:

Low: The health risk of cold injury is Low. Confidence in the health risk estimate is medium (Reference 5, Table 3-6).

9 Noise

9.1 Continuous

No specific hazard sources were documented in the DOEHRS for 2008.

9.1.1 Short- and long-term health risks:

Not evaluated

9.2 Impulse

No specific hazard sources were documented in the DOEHRS for 2008.

9.2.1 Short- and Long-term health risks:

Not evaluated.

10 Unique Incidents/Concerns

10.1 Potential environmental contamination sources

DoD personnel are exposed to various chemical, physical, ergonomic, and biological hazards in the course of performing their mission. These types of hazards depend on the mission of the unit and the operations and tasks which the personnel are required to perform to complete their mission. The health risk associated with these hazards depends on a number of elements including what materials are used, how long the exposure last, what is done to the material, the environment where the task or operation is performed, and what controls are used. The hazards can include exposures to heavy metal particulates (e.g., lead, cadmium, manganese, chromium, and iron oxide), solvents, fuels, oils, and gases (e.g., carbon monoxide, carbon dioxide, oxides of nitrogen, and oxides of sulfur). Most of these exposures occur when performing maintenance task such as painting, grinding, welding, engine repair, or movement through contaminated areas. Exposures to these occupational hazards can occur through inhalation (air), skin contact, or ingestion; however exposures through air are generally associated with the highest health risk.

10.2 Waste Sites/Waste Disposal

No data were available

10.3 Fuel/petroleum products/industrial chemical spills

No data were available.

10.4 Pesticides/Pest Control:

The health risk of exposure to pesticide residues is considered within the framework of typical residential exposure scenarios, based on the types of equipment, techniques, and pesticide products that have been employed, such as enclosed bait stations for rodenticides, various handheld equipment for spot treatments of insecticides and herbicides, and a number of ready-to-use (RTU) methods such as aerosol cans and baits. The control of rodents required the majority of pest management inputs, with the acutely toxic rodenticides staged as solid formulation lethal baits placed in tamper-resistant bait stations indoors and outdoors throughout cantonment areas. Nuisance insects, including biting and stinging insects such as bees, wasps, and ants, also required significant pest management inputs. Use of pesticides targeting against these pests generally involved selection of compounds with low mammalian toxicity and short-term residual using pinpoint rather than broadcast application techniques. No specific hazard sources were documented in the DOEHRS data portal.

10.4.3 Short-term and Long-term health risks

Not evaluated.

10.5 Asbestos

No data was available.

10.6 Lead Based Paint

No data were available.

10.7 Burn Pit

No data were available on burn pits at Camp Panther.

While not specific to Camp Panther, the consolidated epidemiological and environmental sampling and studies on burn pits that have been conducted as of the date of this publication have been unable to determine whether an association does or does not exist between exposures to emissions from the burn pits and long-term health effects (Reference 11). The Institute of Medicine committee's (Reference 11) review of the literature and the data suggests that service in Iraq or Afghanistan (i.e., a broader consideration of air pollution than exposure only to burn pit emissions) may be associated with long-term health effects, particularly in susceptible (e.g., those who have asthma) or highly exposed subpopulations, such as those who worked at the burn pit. Such health effects would be due mainly to high ambient concentrations of PM from both natural and anthropogenic sources, including military sources. If that broader exposure to air pollution turns out to be relevant, potentially related health effects of concern are respiratory and cardiovascular effects and cancer. Susceptibility to the PM health effects could be exacerbated by other exposures, such as stress, smoking, local climatic conditions, and co-exposures to other chemicals that affect the same biologic or chemical processes. Individually, the chemicals measured at burn pit sites in the study were generally below concentrations of health concern for general populations in the United States. However, the possibility of exposure to mixtures of the chemicals raises the potential for health outcomes associated with cumulative exposure to combinations of the constituents of burn pit emissions and emissions from other sources.

11 References

1. Defense Occupational and Environmental Health Readiness System (referred to as the DOEHRSEH database) at <https://doehrs-ih.csd.disa.mil/Doehrs/>.
2. Department of Defense (DoD) Instruction 6490.03, *Deployment Health*, 2006.
3. DoDI 6055.05, Occupational and Environmental Health, 2008.
4. Joint Staff Memorandum (MCM) 0017-12, Procedures for Deployment Health Surveillance, 2012.
5. USAPHC TG230, June 2013 Revision.
6. Modification 9 to United States Central Command Individual Protection and Individual Unit Deployment Policy, 08 September 2008.
7. CDC. 2012. Morbidity and Mortality Weekly Report. Imported Human Rabies in a U.S. Army Soldier. May 4, 2012. 61(17); 302-305.
8. Armed Forces Pest Management Board: <http://www.afpmb.org/content/venomous-animals-country#Afghanistan>. U.S. Army Garrison - Forest Glen, Silver Spring, MD.
9. Clinical Toxinology Resources: <http://www.toxinology.com/>. University of Adelaide, Australia.

10. Goldman RF. 2001. Introduction to heat-related problems in military operations. *In*: Textbook of military medicine: medical aspects of harsh environments Vol. 1, Pandolf KB, and Burr RE (Eds.), Office of the Surgeon General, Department of the Army, Washington DC.
11. IOM (Institute of Medicine). 2011. Long-term health consequences of exposure to burn pits in Iraq and Afghanistan. Washington, DC: The National Academies Press.

12 Where Do I Get More Information?

If a provider feels that the Service member's or Veteran's current medical condition may be attributed to specific OEH exposures at this deployment location, he/she can contact the Service-specific organization below. Organizations external to DoD should contact Deputy Assistant Secretary of Defense for Health Readiness Policy and Oversight (HRP&O).

Army Public Health Center Phone: (800) 222-9698. <http://phc.amedd.army.mil/>

Navy and Marine Corps Public Health Center (NMCPHC) (formerly NEHC) Phone: (757) 953-0700. <http://www.med.navy.mil/sites/nmcphc/Pages/Home.aspx>

U.S. Air Force School of Aerospace Medicine (USAFSAM) (formerly AFIOH) Phone: (888) 232-3764. <http://www.wpafb.af.mil/afri/711hpw/usafsam/>

DoD Health Readiness Policy and Oversight (HRP&O) Phone: (800) 497-6261.
<https://health.mil/Military-Health-Topics/Health-Readiness>